

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 July 2003 (10.07.2003)

PCT

(10) International Publication Number
WO 03/056022 A2(51) International Patent Classification⁷: C12N 15/867,
A01K 67/027(UK) Limited, Medawar Centre, Robert Robinson Avenue,
The Oxford Science Park, Oxford OX4 4GA (GB).

(21) International Application Number: PCT/GB02/05901

(74) Agents: MALLALIEU, Catherine, Louise et al.; D.
Young & Co., 21 New Fetter Lane, London EC4A 1DA
(GB).

(22) International Filing Date:

23 December 2002 (23.12.2002)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0130797.4	21 December 2001 (21.12.2001)	GB
0201140.1	18 January 2002 (18.01.2002)	GB
10/082,122	26 February 2002 (26.02.2002)	US
0211409.8	17 May 2002 (17.05.2002)	GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): OXFORD
BIOMEDICA (UK) LIMITED [GB/GB]; Medawar Cen-
tre, Robert Robinson Avenue, The Oxford Science Park,
Oxford OX4 4GA (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): RADCLIFFE,
Philippa [GB/GB]; c/o Oxford Biomedica (UK) Lim-
ited, Medawar Centre, Robert Robinson Avenue, The
Oxford Science Park, Oxford OX4 4GA (GB). MITRO-
PHANOUS, Kyriacos [GR/GB]; c/o Oxford BioMedica
(UK) Limited, Medawar Centre, Robert Robinson Av-
enue, The Oxford Science Park, Oxford OX4 4GA (GB).
THEMIS, Michael [GB/GB]; c/o Oxford Biomedica

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: TRANSGENIC ORGANISM

(57) Abstract: A method of producing a transgenic cell comprising into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI). Also described is a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising a NOI capable of generating an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA, a micro-RNA or a group I intron. Also described is a viral vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises: (a) a second nucleotide sequence comprising an aptazyme; and (b) a third nucleotide sequence capable of generating a polynucleotide; wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.

- Example 8, page 66

- claims to the main + Fig.

WO 03/056022 A2

Transgenic Organism

Field of the Invention

The present invention relates to a method for producing a transgenic cell and a transgenic organism.

5

Background to the Invention

The ability to introduce genes and/or other DNA sequences into the germline or somatic cells of organisms such as mammals is one of the greatest technical advances in recent biology. Such animals are said to be transgenic. When germline changes are involved, the results of genetic manipulation are inherited by the offspring of the animals and all cells of these offspring inherit the introduced gene and in some cases deleted DNA as part of their genetic make-up. Transgenic mammals have provided a means of studying gene regulation during embryogenesis and in differentiation, for studying the action of oncogenes, and for studying the intricate interactions of cells in the immune system. The whole animal is the ultimate assay system for manipulating genes which direct complex biological processes. In addition, transgenic animals provide exciting possibilities for expressing useful recombinant proteins and for generating precise animal models of human genetic disorders.

20 The production of transgenic animals is commonly done in one of two ways: by targeted insertion of DNA by homologous recombination in embryonic stem (ES) cells which is a labour intensive and time-consuming process, or by pronuclear injection of a fertilised ovum in which integration of DNA is random and may lead to an insertion of large tandem arrays of DNA which are unstable and subject to rearrangements and deletions in subsequent cell divisions. WO99/51755 discusses use of a retroviral expression vector comprising a nucleic acid encoding at least one ribozyme for production of a transgenic animal. No specific disclosure is made of the retrovirus used in the specific example. Mention is also made of the possibility of using an adenovirus, an adeno-associated virus, a lentivirus, a herpes simplex virus or a
30 vaccinia virus. However there are no specific examples of the use of these viruses.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 July 2003 (10.07.2003)

PCT

(10) International Publication Number
WO 2003/056022 A3

(51) International Patent Classification⁷: C12N 15/867, A01K 67/027

(21) International Application Number:
PCT/GB2002/005901

(22) International Filing Date:
23 December 2002 (23.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130797.4 21 December 2001 (21.12.2001) GB
0201140.1 18 January 2002 (18.01.2002) GB
10/082,122 26 February 2002 (26.02.2002) US
0211409.8 17 May 2002 (17.05.2002) GB

(71) Applicant (for all designated States except US): OXFORD BIOMEDICA (UK) LIMITED [GB/GB]; Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).


(72) Inventors; and

(75) Inventors/Applicants (for US only): RADCLIFFE, Philippa [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). MITROPHANOUS, Kyriacos [GR/GB]; c/o Oxford BioMedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). THEMIS, Michael [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).

(74) Agents: MALLALIEU, Catherine, Louise et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

[Continued on next page]

(54) Title: METHOD FOR PRODUCING A TRANSFENIC ORGANISM USING A LENTIVIRAL EXPRESSION VECTOR SUCH AS EIAV



(57) Abstract: A method of producing a transgenic cell comprising into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI). Also described is a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising a NOI capable of generating an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA, a micro-RNA or a group 1 intron. Also described is a viral vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises: (a) a second nucleotide sequence comprising an aptazyme; and (b) a third nucleotide sequence capable of generating a polynucleotide; wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.

WO 2003/056022 A3



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,

TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report

(88) **Date of publication of the international search report:**
31 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PL 17 GB 02/05901

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/867 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE, MEDLINE, PASCAL, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 312 683 B1 (CARROLL MILES WILLIAM ET AL) 6 November 2001 (2001-11-06) examples 4,5	1-6,15, 32,62
X	YAMADA K ET AL: "Functional correction of fanconi anemia group C hematopoietic cells by the use of a novel lentiviral vector." MOLECULAR THERAPY: THE JOURNAL OF THE AMERICAN SOCIETY OF GENE THERAPY. UNITED STATES APR 2001, vol. 3, no. 4, April 2001 (2001-04), pages 485-490, XP001147753 ISSN: 1525-0016 page 490, left-hand column, last paragraph; figure 2	1-6,15, 32,62

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

17 April 2003

Date of mailing of the international search report

10.09.03

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lanzrein, M

INTERNATIONAL SEARCH REPORT

Inte pnal Application No
PC 1, GB 02/05901

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TARANTAL A F ET AL: "Lentiviral vector gene transfer into fetal rhesus monkeys (Macaca mulatta): lung-targeting approaches."</p> <p>MOLECULAR THERAPY: THE JOURNAL OF THE AMERICAN SOCIETY OF GENE THERAPY. UNITED STATES DEC 2001, vol. 4, no. 6, December 2001 (2001-12), pages 614-621, XP002238838</p> <p>ISSN: 1525-0016</p> <p>figures 1,3,5; table 2</p> <p>-----</p>	<p>1-16, 32-39, 62-65</p>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 02/05901

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-16, 62 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-16 (completely); 32-39, 62-65 (partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 02 05901

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-16 (completely); 32-39, 62-65 (partially)

A method for producing a transgenic cell using a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI); the transgenic cell produced by said method; a transgenic organism which is generated from said transgenic cell.

2. Claims: 17-31 (completely); 32-39, 62-65 (partially)

A method for producing a transgenic cell by introducing into a non-dividing cell lentiviral expression vector comprising a nucleotide of interest (NOI); the transgenic cell produced by said method; a transgenic organism which is generated from said transgenic cell.

3. Claims: 40-61 (completely); 62-65 (partially)

A vector comprising an aptazyme operably linked to a sequence capable of generating a polynucleotide, wherein the aptazyme is activable to cleave a transcript such that said polynucleotide is generated; method for producing a transgenic cell using said vector; a transgenic organism generated from said transgenic cell.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 02/05901

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6312683	B1	06-11-2001	
		AU 760287 B2	08-05-2003
		AU 1773299 A	12-07-1999
		CA 2315269 A1	01-07-1999
		CN 1285000 T	21-02-2001
		EP 1042493 A1	11-10-2000
		GB 2356200 A ,B	16-05-2001
		WO 9932646 A1	01-07-1999
		GB 2334257 A ,B	18-08-1999
		GB 2345062 A ,B	28-06-2000
		JP 2002516065 T	04-06-2002

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 July 2003 (10.07.2003)

PCT

(10) International Publication Number
WO 2003/056022 A3

(51) International Patent Classification⁷: C12N 15/867, A01K 67/027

(21) International Application Number:
PCT/GB2002/005901

(22) International Filing Date:
23 December 2002 (23.12.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0130797.4 21 December 2001 (21.12.2001) GB
0201140.1 18 January 2002 (18.01.2002) GB
10/082,122 26 February 2002 (26.02.2002) US
0211409.8 17 May 2002 (17.05.2002) GB

(71) Applicant (for all designated States except US): OXFORD BIOMEDICA (UK) LIMITED [GB/GB]; Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).

(72) Inventors; and


(75) Inventors/Applicants (for US only): RADCLIFFE, Philippa [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). MITROPHANOUS, Kyriacos [GR/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). THEMIS, Michael [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).

(74) Agents: MALLALIEU, Catherine, Louise et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

[Continued on next page]

(54) Title: METHOD FOR PRODUCING A TRANSGENIC ORGANISM USING A LENTIVIRAL EXPRESSION VECTOR SUCH AS EIAV

Schematic of pONY8.1Zhyb



(57) Abstract: A method of producing a transgenic cell comprising introducing into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI). Also described is a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising a NOI capable of generating an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA, a micro-RNA or a group 1 intron. Also described is a viral vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises: (a) a second nucleotide sequence comprising an aptazyme; and (b) a third nucleotide sequence capable of generating a polynucleotide; wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *with international search report*

— *with amended claims*

(88) **Date of publication of the international search report:**
31 December 2003

Date of publication of the amended claims: 12 February 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

AMENDED CLAIMS

[received by the International Bureau on 10 November 2003 (10.11.03);
new claims 66-70 added; remaining claims unchanged (1 page)]

66. A method of producing a transgenic cell comprising introducing into a cell a viral expression vector comprising a nucleotide of interest (NOI) which encodes and is capable of expressing a therapeutic protein.

5

67. A method of producing a transgenic cell comprising introducing into a cell a viral expression vector comprising an aptazyme capable of generating an siRNA, an antisense oligonucleotide, a ribozyme, a short hairpin RNA, a micro-RNA or a group I intron.

10

68. A transgenic cell produced by the method of claim 66 or 67.

69. A transgenic organism which is generated from or obtainable by generation from a transgenic cell according to claim 68, or from the method as defined in claim 66 or 67.

15

70. A transgenic egg derived from a transgenic organism according to claim 69.

Thus, in recent years, retroviruses have been proposed for use in gene therapy. Essentially, retroviruses are RNA viruses with a life cycle different to that of lytic viruses. In this regard, when a retrovirus infects a cell, its genome is converted to a DNA form. In other words, a retrovirus is an infectious entity that replicates through a DNA intermediate. More details on retroviral infection etc. are presented later on.

There are many retroviruses and examples include: murine leukaemia virus (MLV), mouse mammary tumour virus (MMTV), Rous sarcoma virus (RSV), Fujinami sarcoma virus (FuSV), Moloney murine leukemia virus (Mo-MLV), FBR murine osteosarcoma virus (FBR MSV), Moloney murine sarcoma virus (Mo-MSV), Abelson murine leukemia virus (A-MLV), Avian myelocytomatosis virus-29 (MC29), and Avian erythroblastosis virus (AEV).

There is also a family called lentiviruses including human immunodeficiency virus (HIV) and equine infectious anaemia virus (EIAV). Further details are given below.

A detailed list of retroviruses and lentiviruses may be found in Coffin *et al* ("Retroviruses" 1997 Cold Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763).

Details on the genomic structure of some retroviruses may be found in the art. By way of example, details on HIV may be found from the NCBI Genbank (i.e. Genome Accession No. AF033819).

As indicated above, there has been considerable interest in the development of retroviral vector systems based on lentiviruses, a small subgroup of the retroviruses. This interest arises firstly from the notion of using HIV-based vectors to target anti-HIV therapeutic genes to HIV susceptible cells and secondly from the prediction that, because lentiviruses are able to infect non-dividing cells (Lewis & Emerman 1993 J.Virol. 68, 510), vector systems based on these viruses would be able to transduce non-dividing cells (e.g. Vile & Russel 1995 Brit. Med. Bull. 51, 12). Vector systems based on HIV have been produced (Buchsacher & Panganiban 1992 J.Virol. 66, 2731) and they have been used to

transduce CD4+ cells and, as anticipated, non-dividing cells (Naldini *et al*, 1996 Science 272, 263). In addition lentiviral vectors enable very stable long-term expression of the gene of interest. This has been shown to be at least three months for transduced rat neuronal cells. The MLV based vectors were only able to express the gene of interest
5 for six weeks.

HIV-based vectors produced to date result in an integrated provirus in the transduced cell that has HIV LTRs at its ends. This limits the use of these vectors as the LTRs have to be used as expression signals for any inserted gene unless an internal promoter is used. The
10 use of internal promoters has significant disadvantages. The unpredictable outcome of placing additional promoters within the retroviral LTR transcription unit is well documented (Bowtell *et al*, 1988 J.Virol. 62, 2464; Correll *et al*, 1994 Blood 84, 1812; Emerman and Temin 1984 Cell 39, 459; Ghattas *et al*, 1991 Mol.Cell.Biol. 11, 5848; Hantzopoulos *et al*, 1989 PNAS 86, 3519; Hatzoglou *et al*, 1991 J.Biol.Chem 266, 8416;
15 Hatzoglou *et al*, 1988 J.Biol.Chem 263, 17798; Li *et al*, 1992 Hum.Gen.Ther. 3, 381; McLachlin *et al*, 1993 Virol. 195, 1; Overell *et al*, 1988 Mol.Cell Biol. 8, 1803; Scharfman *et al*, 1991 PNAS 88, 4626; Vile *et al*, 1994 Gene Ther 1, 307; Xu *et al*, 1989 Virol. 171, 331; Yee *et al*, 1987 PNAS 84, 5197). The factors involved appear to include the relative position and orientation of the two promoters, the nature of the promoters and
20 the expressed genes and any selection procedures that may be adopted. The presence of internal promoters can affect both the transduction titers attainable from a packaging cell line and the stability of the integrated vector.

HIV and other lentiviral LTRs have virus-specific requirements for gene expression. For
25 example, the HIV LTR is not active in the absence of the viral Tat protein (Cullen 1995 AIDS 9, S19). It is desirable, therefore, to modify the LTRs in such a way as to change the requirements for gene expression. In particular tissue specific gene expression signals may be required for some gene therapy applications.

30 HIV vectors have a number of significant disadvantages which may limit their therapeutic application to certain diseases. HIV-1 has the disadvantage of being a human pathogen carrying potentially oncogenic proteins and sequences. There is the

risk that introduction of vector particles produced in packaging cells which express HIV gag-pol will introduce these proteins into the patient leading to seroconversion. For these reasons, there is a need to develop lentiviral-based vectors which do not introduce HIV proteins into patients.

5

The use of MLV in the production of transgenic animals has also been proposed. However, the level of expression using this technique has been disappointing.

There is also a long felt need to generate disease models in which the transgene is regulatable and which can be generated efficiently.

10

Aspects of the present invention overcome these problems.

According to a first aspect of the present invention there is provided a method of producing a transgenic cell comprising introducing into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI).

15

The present invention provides an efficient way of producing transgenic animals and which overcomes any potential difficulties associated with the use of primate lentiviruses.

20

Preferably, the non-primate lentiviral expression vector is derived from ELAV, FIV, BIV, CAEV or MVV, with ELAV being particularly preferred.

One of the advantages of the present invention is that the expression vector can be introduced in vivo or ex vivo. In one embodiment the method is carried out in vitro. In another embodiment, the cell is in utero.

25

Several methods for introducing foreign DNA into the germline of mammals have been developed. The techniques allow the mixing of cells from different embryos, i.e. chimaera production, introducing pluripotent cells such as ES cells into developing embryos, micro-injecting DNA, and infection by retroviruses. Many of these

30

techniques have the fundamental requirement of removing fertilised eggs or early embryos, culturing them in vitro and then returning them to foster mothers where further embryogenesis can proceed. In particular the production of transgenic animals by targeted insertion of DNA by homologous recombination in ES cells is a labour
5 intensive and time-consuming process with, e.g. a turnaround time of 8 to 9 weeks from nuclear injection.

One major advantage of this embodiment of the present invention is the ability to avoid the need to remove, culture in vitro and then reimplant cells. It also avoids the
10 intensive and time-consuming production of recombinant ES cells.

Indeed, a vast number of genes of unknown function are now available following large scale gene sequencing programmes. To develop therapeutic products from novel genomic targets, it will be necessary to correlate biology with gene sequence
15 information. The present invention provides an efficient and effective in vivo method for assisting in the validation of targets.

We have also found that good levels of expression may be achieved using the present invention, and that the levels of expression, particularly with ELAV, are better than
20 those achieved using MLV. This is surprising.

Another advantage of the present invention is its efficiency. Regulatable knock-out disease models can be efficiently produced though transduction with one vector, if desired, and few generations are required. Thus, the present invention meets a long
25 felt want whose solution was not obvious at the time.

A further advantage of this aspect of the present invention is its flexibility; the lentiviral vector can be introduced throughout the development of the organism. Thus in one embodiment the cell is a perinatal cell, which could be an embryonic cell. In a
30 particular aspect of this embodiment the embryonic cell is in utero. However, the method may be applied to any cell such as any somatic cell and also any cell which is capable of giving rise to a germ line change. Such cells include the germ cells, of

course, but the present invention can also be applied to a cell which is involved either directly or indirectly in gametogenesis or fertilisation. We also include equivalent cells which are arrived at without direct fertilisation, e.g. through cell nuclear replacement techniques.

5

Preferably the cell is an oocyte, an oviduct cell, an ovarian cell, an ovum, an ES cell, a blastocyte, a spermatocyte, a spermatid, a spermatozoa, or a spermatogonia.

When seeking to achieve germ line changes, it will be appreciated that the earlier
10 transduction occurs, the better as there is a greater chance of transducing germ cells.

A particular advantage with the use of lentiviral vectors is that it is possible to transduce non- or slowly- dividing cells, such as oocytes and sperm-forming cells.

15 Therefore according to another aspect of the present invention there is provided a method for producing a transgenic cell comprising introducing into a nondividing cell a lentiviral expression vector comprising an NOI. The lentiviral expression vector may be derived from a non-primate lentivirus, but may also be derived from a primate lentivirus such as HIV.

20

An important advantage of this aspect of the present invention is that cells do not have to be fertilised for transduction to be achieved.

By non-dividing cells we include cells which are capable of dividing but are non-
25 dividing at a particular time.

The method is not limited to a particular cell type, but the cell is preferably a eukaryotic cell, such as an animal, preferably mammalian, or yeast cell. Examples of cells to which the present invention is applicable include murine, human, porcine,
30 bovine, simian, ovine, equine, avian such as fowl, particularly chickens, insect or reptile or piscine cell. The cell may be from, e.g., *C. elegans* or *drosophila*.

In one embodiment, the cell is from a non-human organism.

Preferably the lentiviral expression vector is pseudotyped.

- 5 Preferably the lentiviral expression vector does not contain any functional accessory genes.

The NOI may be operably linked to a constitutive, tissue-specific or an inducible promoter.

10

Preferably, the NOI encodes and is capable of expressing a therapeutic protein, or encodes an antisense oligonucleotide or encodes a ribozyme.

- 15 In a preferred embodiment the NOI is capable of generating an RNA molecule capable of post-transcriptional silencing of a target gene. In this aspect of the present invention we provide a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising an NOI capable of generating an RNA molecule capable of post-transcriptional silencing of a target gene.

- 20 Preferably, the NOI is capable of generating a short RNA, a siRNA, a short hairpin RNA, a micro-RNA or a group I intron.

- 25 In one embodiment, expression of a short RNA, a siRNA, a short hairpin RNA, a micro-RNA is regulated by a tetracycline-responsive derivative of an RNA polymerase promoter.

- 30 In one embodiment the method comprising introducing at least one NOI which is capable of expressing a protein, preferably a therapeutic protein, and at least one NOI which is capable of generating an RNA molecule capable of post-transcriptional silencing of a target gene. The at least one NOI which is capable of expressing a protein, preferably a therapeutic protein, and the at least one NOI which is capable of

generating an RNA molecule capable of post-transcriptional silencing of a target gene may be incorporated within the same or separate lentiviral expression vectors.

5 The lentiviral expression vector may be introduced into a target cell through administration via any convenient route of access, such as a cell of the umbilical cord, placenta, or amniotic fluid; or directly into an organ such as the uterus, gonad, brain, kidney, liver, heart, bone marrow, blood, central nervous system, or lung.

10 One problem associated with the production of transgenic animals for establishing disease models arises where the loss of expression in say a knock out mouse is lethal. In the methods of the present invention the NOI can be operably linked to a tissue-specific or an inducible promoter. This is particularly advantageous where ablation of gene expression is desired at a particular developmental stage or in a specific tissue.

15 The NOI may be expressed in the transgenic organism in a constitutive, tissue-specific or regulatable manner. Examples of cells where the NOI may be expressed include a cell of any organ or tissue, such as a cell of the brain, kidney, liver, heart, bone marrow, blood, central nervous system, or lung of said organism. The NOI may also be expressed at a particular developmental stage of the organism.

20 As discussed above, in a preferred embodiment of the present invention, said NOI encodes an RNA e.g., a short RNA, a siRNA, a short hairpin RNA or a micro-RNA capable of post-transcriptional silencing of a target gene.

25 The present invention also relates to transgenic animals derivable from such cells. As well as their use in disease models it will be appreciated that such transgenic animals can be used in the production of proteins, such a therapeutic proteins, e.g. insulin.

30 In a particular aspect of the present invention there is provided an egg derived from a transgenic avian. Thus according to this aspect of the present invention there is provided a method for generating a transgenic avian and/or egg comprising introducing into an avian cell a lentiviral expression vector comprising an NOI. In the

case where the NOI encodes a protein, the protein can be cleanly and efficiently harvested from the transgenic egg expressing the protein.

However, we have recognised that there is a limit on the volume of protein which can be produced in an egg due to its size. We have now found that by silencing one or more of the genes encoding proteins which are naturally present in the egg it is possible to increase the yield of the protein expressed by the introduced NOI. These natural proteins may be down-regulated through the method of the present invention, i.e. through the use of a lentiviral expression vector encoding at least one NOI which is capable of generating an RNA molecule capable of post-transcriptional silencing of a natural egg gene. The at least one NOI which is capable of expressing a protein, preferably a therapeutic protein, and the at least one NOI which is capable of generating an RNA molecule capable of post-transcriptional silencing of the target egg gene may be incorporated within the same or separate lentiviral expression vectors. In a preferred embodiment of this aspect of the present invention the NOI capable of expressing the protein is placed under the control of a promoter which is native to the egg, such as the lysozyme promoter.

Thus the present invention provides a transgenic organism or egg comprising at least one NOI capable of expressing a protein, preferably a therapeutic protein, and at least one NOI which is capable of generating an RNA molecule capable of post-transcriptional silencing of a natural egg gene. As discussed above this second NOI may encode an RNA e.g., a short RNA, a siRNA, a short hairpin RNA or a micro-RNA capable of post-transcriptional silencing of a target gene.

25

Post-transcriptional gene silencing (PTGS) mediated by double-stranded dsRNA is a conserved cellular defence mechanism for controlling the expression of foreign genes. It is thought that the random integration of elements such as transposons or viruses causes the expression of dsRNA which activates sequence-specific degradation of homologous single-stranded mRNA or viral genomic RNA. The silencing effect is known as RNA interference (RNAi). The mechanism of RNAi involves the processing of long dsRNAs into duplexes of 21-25 nucleotide (nt) RNAs. These products are

30

called small interfering or silencing RNAs (siRNAs) which are the sequence-specific mediators of mRNA degradation. In addition to siRNAs, the expression of short RNAs may act to redirect splicing ('exon-skipping') or polyadenylation or to inhibit translation.

5

Although viral vectors are efficient tools for in vivo gene delivery, the short length of RNA molecules involved in post-transcriptional gene silencing means that transcription of these RNAs using conventional expression cassettes is difficult. A further problem is that the use of viral vectors, e.g., lentiviral vectors, for generating transgenics to deliver the aforementioned RNA molecules which target a gene product with an important or essential function may result in death of the transgenic animal during development. An aspect of the present invention overcomes this problem by providing vectors in which transcription of polynucleotides e.g., siRNAs, are able to be regulated by use of an aptazyme. Indeed this aspect of the invention is not limited to methods involving lentiviral vectors and constitutes an independent aspect of the invention.

10
15

Thus, in a second aspect of the invention there is provided a vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises:

20

- (a) a second nucleotide sequence encoding an aptazyme; and
- (b) a third nucleotide sequence capable of generating a polynucleotide;

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.

25

In other words there is provided a vector comprising a nucleic acid sequence, wherein said nucleic acid sequence comprises:

30

- (c) a first nucleotide sequence encoding an aptazyme; and
- (d) a second nucleotide sequence capable of generating a polynucleotide;

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the nucleic acid sequence such that said polynucleotide is generated.

5

Preferably the vector is a viral vector.

In a preferred embodiment of this aspect of the invention, the polynucleotide is an RNA molecule capable of modulating expression of a target gene. Preferably, the
10 RNA molecule is selected from the group comprising siRNA, short hairpin RNA, microRNA, anti-sense RNA and a ribozyme.

Aptazymes are allosteric ribozymes. Aptamers are nucleic acid molecules which form structures which are able to bind a number of ligands including proteins and drug
15 molecules. By replacing one helix of a ribozyme, e.g. a hammerhead ribozyme, with an aptamer it has been possible to create a catalytic RNA which is able to cleave a substrate (which may be itself) as the result of conformational change induced by the presence or absence of a ligand.. Aptazymes which can be induced or inhibited by flavin mononucleotide (FMN) have been described (Soukup and Breaker 1999), as has
20 an aptazyme which is inhibited by doxycycline (Piganaeu et al 2000).

In a further independent aspect of the present invention, cleavage induced by an aptazyme may be used to directly modulate expression of a NOI.

25 Thus, according to a third aspect of the present invention there is provided a vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises:

- (a) a second nucleotide sequence encoding an aptazyme; and
- 30 (b) a third nucleotide sequence comprising a NOI;

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave the transcript of the first nucleotide sequence such that expression of said NOI is inhibited.

- 5 In other words the present invention provides a vector comprising a nucleic acid sequence, wherein said nucleic acid sequence comprises:

- (c) a first nucleotide sequence encoding an aptazyme; and
- (d) a second nucleotide sequence comprising a NOI;

10

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave the transcript of the nucleic acid sequence such that expression of said NOI is inhibited.

- 15 Preferably the vector is a viral vector.

In one embodiment the aptazyme encoded by the above vector is activatable to cleave the transcript of the first nucleotide sequence at a position within the transcript of the third nucleotide sequence.

20

In a preferred embodiment the NOI encoded by the vector according to the third aspect of the present invention encodes a therapeutic protein.

- 25 In one embodiment of the second and third aspects of the present invention the aptazyme is activated by binding of a ligand to the aptazyme.

In another embodiment of the second and third aspects of the present invention the aptazyme is deactivated by binding of a ligand to the aptazyme.

- 30 In a preferred embodiment of the second and third aspects of the present invention the vector further comprises a fourth nucleotide sequence encoding a ligand capable of

binding the aptazyme. The nucleotide sequence encoding the ligand may be operatively linked to a promoter.

The ligand may be selected from the group comprising polypeptides and fragments thereof, linear peptides, cyclic peptides, and nucleic acids which encode therefor, synthetic and natural compounds including low molecular weight organic or inorganic compounds and antibodies.

- 5 In preferred embodiments the ligand for use in these aspects of the invention is selected from the group comprising FMN, doxycycline and VEGF, tetracycline or glucose.

- 10 In another embodiment of the second and third aspects of the present invention (a) and (b), are operably linked to a promoter. Preferred promoters are selected from the group comprising RNA polymerase III (U6) promoters such as the U6 promoter as well as conventional RNA polymerase II promoters.

- 15 The promoter may be operably linked to at least one copy of a tetracycline responsive element (TRE), e.g., the Tet operator, such that transcription of the first nucleotide sequence is regulated by a tetracycline modulator and tetracycline or derivatives thereof.

- 20 In one embodiment, the vector according to the second and third aspects of the present invention comprises a fifth nucleotide sequence encoding a tetracycline modulator.

- 25 Although the production of the vector is preferably carried out under conditions which should minimise activity of the aptazyme, and hence unwanted destruction of the vector genome by self-cleavage, in a preferred embodiment the vector is configured as a split intron vector. This ensures that the full sequence of the aptazyme is only present in the transcript encoded by the provirus and not in the RNA genome present in the vector particle. An additional means of preventing formation of a potentially active aptazyme within the viral RNA genome is to use a promoter containing a sequence at

its 3' end which is able to base-pair with a part of the aptazyme such as to form a hairpin to prevent formation of active aptazyme within the viral RNA genome. Details of split intron vectors are described in WO 99/15683.

- 5 The vector according to the second and third aspects of the present invention may be derived from any suitable virus, for example a retrovirus, a lentivirus, an adenovirus, an adeno-associated vector, a herpes vector, a pox viral vector, a parvovirus vector or a baculoviral vector.
- 10 In a fourth aspect of the present invention there is provided a method of producing a transgenic cell using a vector of the second or third aspects of the present invention

According to a fifth aspect of the present invention there is provided a transgenic cell produced according to any of the methods of the present invention.

15

According to a sixth aspect of the present invention there is provided a transgenic organism which is generated from or obtainable by generation from a transgenic cell of the present invention.

- 20 According to a seventh aspect of the present invention there is provided a transgenic organism of the present invention wherein the NOI is expressed in a haematopoietic cell, (including monocytes, macrophages, lymphocytes, granulocytes, or progenitor cells of any of these); endothelial cell, tumour cell, stromal cell, astrocyte, or glial cell, muscle cell, epithelial cell, neuron, fibroblast, hepatocyte. astrocyte, kidney, liver,
- 25 heart or lung cell.

- In another aspect of the present invention there is provided a transgenic organism according to the present invention wherein the NOI is expressed in an oviduct cell, reproductive tract cell, albumin, haematopoietic cell, (including monocytes,
- 30 macrophages, lymphocytes, granulocytes, or progenitor cells of any of these); endothelial cell, tumour cell, stromal cell, astrocyte, or glial cell, muscle cell, epithelial cell, neuron, fibroblast, hepatocyte. astrocyte, kidney, liver, heart or lung cell.

Various preferred features and embodiments of the present invention will now be described by way of non-limiting example and with reference to the accompanying drawings in which:

5

Description of the Figures

Figure 1 shows a liver and tissue histology after in utero injection of EIAV lentivirus and shows sections of mouse liver stained for the β -galactosidase marker gene 3, 7, 14, 28, 79 days and 6 months after foetal intravenous injection;

- 10 Figure 2 shows tissue and histology after in utero injection of EIAV lentivirus and shows section of mouse liver, heart, skeletal muscle, lung, brain, and kidney stained for the β -galactosidase marker gene at variously 3, 7, 14 and 79 days after foetal intravenous injection;

- 15 Figure 3 shows a mouse dorsal root ganglia stained for the β -galactosidase marker gene 7 days post foetal intraspinal injection of EIAV viral vector expressing nuclear localising LacZ;

Figure 4 shows a section of a mouse dorsal root ganglia stained for the β -galactosidase marker gene days post foetal intraspinal injection of EIAV viral vector expressing nuclear localising LacZ;

- 20 Figure 5 shows a section of mouse liver stained for the β -galactosidase marker gene 7 days post foetal intravenous injection of EIAV viral vector expression nuclear localising LacZ;

- 25 Figure 6 shows a mouse renal glomeruli and a section thereof stained for the β -galactosidase marker gene 7 days post foetal intravenous injection of EIAV viral vector expression nuclear localising LacZ;

Figure 7 shows a mouse pancreas and a section thereof stained for the β -galactosidase marker gene 7 days post foetal intravenous injection of EIAV viral vector expression nuclear localising LacZ;

Figure 8 shows mouse skeletal muscle stained for the β -galactosidase marker gene 7 days post foetal intramuscular injection of EIAV viral vector expression nuclear localising LacZ;

Figure 9 shows a mouse diaphragm and planar and transverse sections thereof stained
5 for the β -galactosidase marker gene two weeks post foetal intraperitoneal injection of EIAV viral vector expression LacZ;

Figure 10 shows a mouse leg and planar and transverse sections thereof stained for the β -galactosidase marker gene two weeks post foetal intramuscular injection of EIAV viral vector expression nuclear localising LacZ;

10 Figure 11 shows X-Gal visualisation for β -galactosidase 96 hours after intra-thoracic and intra-peritoneal injection of EIAV viral vector. Figure 11A shows a sagittal section with the viscera removed. The diaphragm has been excised and is viewed anteriorly in Figure 11B;

Figure 12 shows schematic representations of EIAV genomes with sizes. These may
15 be used for transfection in the present invention. Upon transfection the 3' LTR will be copied to the 5' LTR;

Figure 13 shows the nucleic acid sequence of pONY8.1G;

Figure 14 shows the nucleic acid sequence of pONY8.4ZCG;

Figure 15 shows the nucleic acid sequence of pONY8.4GCZ;

20 Figure 16 is a schematic representation of the hybrid U3 region of a vector for use in the present invention;

Figure 17 shows the nucleic acid sequence of this hybrid LTR;

Figure 18 shows the nucleic acid sequence of pONY8.1ZHyb;

Figure 19 is a schematic representation of pONY8.1ZHyb;

25 Figures 20 (a)-(c) show expression cassettes for use in RNAi applications;

Figure 21 (a) shows an expression cassette for use in mediating aptazyme regulated siRNA gene silencing;

Figure 21 (b) shows the structure of the transcripts of the expression cassette of Fig 24a.

Figure 22 shows an expression cassette for use in hypoxically inducing silencing of VEGF by siRNAs;

- 5 Figure 23 (a) shows an expression cassette comprising an RNA polymerase II promoter for expressing aptazyme regulated short hairpin;

Figure 23 (b) shows an expression cassette comprising an RNA polymerase II promoter for expressing aptazyme regulated antisense siRNA;

- 10 Figure 24 (a) shows an expression cassette for use in mediating aptazyme regulated insulin expression;

Figure 24 (b) shows an expression cassette for use in mediating aptazyme regulated Factor IX expression;

Figure 25 (a) shows a schematic of a split intron strategy to avoid self-cleavage of RNA genome;

- 15 Figure 25 (b) shows an expression cassette for use in a split intron strategy;

Figure 26 shows a schematic of a double hairpin strategy to avoid self-cleavage of RNA genome.

Detailed Description of the Invention

- 20 Although in general the techniques mentioned herein are well known in the art, reference may be made in particular to Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (1989) and Ausubel *et al.*, Short Protocols in Molecular Biology (1999) 4th Ed, John Wiley & Sons, Inc.

- 25 One aspect of the present invention relates to a method of producing a transgenic cell using a non-primate lentiviral expression vector and a transgenic organism which is obtainable from the transgenic cell or of which the transgenic cell forms part. More particularly, this aspect of the present invention relates to a lentiviral vector useful in

gene therapy and in the production of disease models. The development of disease models, e.g. transgenic "knockout" mice, has greatly benefited studies of gene function, with particular relevance in establishing mammalian models of genetic disease.

5

Gene therapy includes any one or more of: the addition, the replacement, the deletion, the supplementation, the manipulation etc. of one or more nucleotide sequences in, for example, one or more targeted sites - such as targeted cells. If the targeted sites are targeted cells, then the cells may be part of a tissue or an organ. General teachings on gene therapy may be found in Molecular Biology (Ed Robert Meyers, Pub VCH, such as pages 556-558).

By way of further example, gene therapy also provides a means by which any one or more of: a nucleotide sequence, such as a gene, can be applied to replace or supplement a defective gene; a pathogenic gene or gene product can be eliminated; a new gene can be added in order, for example, to create a more favourable phenotype; cells can be manipulated at the molecular level to treat cancer (Schmidt-Wolf and Schmidt-Wolf, 1994, Annals of Hematology 69:273-279) or other conditions - such as immune, cardiovascular, neurological, inflammatory or infectious disorders; antigens can be manipulated and/or introduced to elicit an immune response - such as genetic vaccination.

A transgenic organism is an organism which includes in at least one of its cells a nucleotide of interest (NOI). In one embodiment the cell is a germline cell. In another embodiment, the cell is a somatic cell. More particularly, the NOI has been introduced experimentally, e.g. using cDNA technology.

The NOI is commonly referred to as a "transgene", i.e. a gene that is inserted into the cell in such a way that ensures its function. When the gene is inserted into a germ line gene it should function, replicate and be transmitted as a normal gene.

The present invention encompasses chimeras and mosaics.

A "chimera" is an organism composed of a mixture of genetically different cells.

5 A "mosaic" is an organism in which the transgene is incorporated into the genome after the first cell division. The organism will be mosaic as different cells will have different sites of integration.

10 A transgenic organism of the invention is preferably a multicellular eukaryotic organism, such as an animal or a plant, or a fungus, or a unicellular eukaryotic organism such as a yeast.

Then organism is preferably an animal, more preferably a mammal.

15 The first aspect of the present invention employs a non-primate lentiviral expression vector.

20 As it is well known in the art, a vector is a tool that allows or facilitates the transfer of an entity from one environment to another. In accordance with the present invention, and by way of example, some vectors used in recombinant DNA techniques allow entities, such as a segment of DNA (such as a heterologous DNA segment, such as a heterologous cDNA segment), to be transferred into a host cell for the purpose of replicating the vectors comprising a segment of DNA. Examples of vectors used in recombinant DNA techniques include but are not limited to plasmids, chromosomes, artificial chromosomes or viruses.

25

The term "expression vector" means a construct capable of *in vivo* or *in vitro/ex vivo* expression.

30 The lentiviral vector used in aspects of the present invention is capable of transducing a target non-dividing cell. One advantage of this feature is that since freshly isolated

oocytes are quiescent, transduction rates may be enhanced by the use of lentiviral rather than retroviral vectors.

5 In a typical vector for use in the method of the present invention, at least part of one or more protein coding regions essential for replication may be removed from the virus. This makes the viral vector replication-defective. Portions of the viral genome may also be replaced by a library encoding candidate modulating moieties operably linked to a regulatory control region and a reporter moiety in the vector genome in order to generate a vector comprising candidate modulating moieties which is capable of
10 transducing a target non-dividing host cell and/or integrating its genome into a host genome.

Preferably the viral vector capable of transducing a target non-dividing or slowly dividing cell is a lentiviral vector.

15

Lentivirus vectors are part of a larger group of retroviral vectors. A detailed list of lentiviruses may be found in Coffin *et al* ("Retroviruses" 1997 Cold Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763). In brief, lentiviruses can be divided into primate and non-primate groups. Examples of primate
20 lentiviruses include but are not limited to: the human immunodeficiency virus (HIV), the causative agent of human auto-immunodeficiency syndrome (AIDS), and the simian immunodeficiency virus (SIV). The non-primate lentiviral group includes the prototype "slow virus" visna/maedi virus (VMV), as well as the related caprine arthritis-encephalitis virus (CAEV), equine infectious anaemia virus (EIAV) and the
25 more recently described feline immunodeficiency virus (FIV) and bovine immunodeficiency virus (BIV).

A distinction between the lentivirus family and other types of retroviruses is that lentiviruses have the capability to infect both dividing and non-dividing cells (Lewis *et al* 1992 EMBO. J 11: 3053-3058; Lewis and Emerman 1994 J. Virol. 68: 510-516). In
30 contrast, other retroviruses - such as MLV - are unable to infect non-dividing or slowly

dividing cells such as those that make up, for example, muscle, brain, lung and liver tissue.

5 A "non-primate" vector, as used herein in some aspects of the present invention, refers to a vector derived from a virus which does not primarily infect primates, especially humans. Thus, non-primate virus vectors include vectors which infect non-primate mammals, such as dogs, sheep and horses, reptiles, birds and insects.

10 A lentiviral or lentivirus vector, as used herein, is a vector which comprises at least one component part derivable from a lentivirus. Preferably, that component part is involved in the biological mechanisms by which the vector infects cells, expresses genes or is replicated. The term "derivable" is used in its normal sense as meaning the sequence need not necessarily be obtained from a retrovirus but instead could be derived therefrom. By way of example, the sequence may be prepared synthetically or by use of
15 recombinant DNA techniques.

The non-primate lentivirus may be any member of the family of lentiviridae which does not naturally infect a primate and may include a feline immunodeficiency virus (FIV), a bovine immunodeficiency virus (BIV), a caprine arthritis encephalitis virus (CAEV), a Maedi visna virus (MVV) or an equine infectious anaemia virus (EIAV).
20 Preferably the lentivirus is an EIAV. Equine infectious anaemia virus infects all equidae resulting in plasma viremia and thrombocytopenia (Clabough, et al. 1991. J Virol. 65:6242-51). Virus replication is thought to be controlled by the process of maturation of monocytes into macrophages.

25

In one embodiment the viral vector is derived from EIAV. EIAV has the simplest genomic structure of the lentiviruses and is particularly preferred for use in the present invention. In addition to the *gag*, *pol* and *env* genes EIAV encodes three other genes: *tat*, *rev*, and *S2*. *Tat* acts as a transcriptional activator of the viral LTR (Derse and Newbold 1993 Virology. 194:530-6; Maury, et al 1994 Virology. 200:632-42) and *Rev* regulates and coordinates the expression of viral genes through *rev*-response elements (RRE) (Martarano et al 1994 J Virol. 68:3102-11). The mechanisms of action of these
30

two proteins are thought to be broadly similar to the analogous mechanisms in the primate viruses (Martano et al ibid). The function of S2 is unknown. In addition, an EIAV protein, Ttm, has been identified that is encoded by the first exon of *tat* spliced to the *env* coding sequence at the start of the transmembrane protein.

5

In addition to protease, reverse transcriptase and integrase non-primate lentiviruses contain a fourth *pol* gene product which codes for a dUTPase. This may play a role in the ability of these lentiviruses to infect certain non-dividing cell types.

- 10 The viral RNA of this aspect of the invention is transcribed from a promoter, which may be of viral or non-viral origin, but which is capable of directing expression in a eukaryotic cell such as a mammalian cell. Optionally an enhancer is added, either upstream of the promoter or downstream. The RNA transcript is terminated at a polyadenylation site which may be the one provided in the lentiviral 3' LTR or a
15 different polyadenylation signal.

Thus the present invention employs a DNA transcription unit comprising a promoter and optionally an enhancer capable of directing expression of a non-primate lentiviral vector genome.

20

- Transcription units as described herein comprise regions of nucleic acid containing sequences capable of being transcribed. Thus, sequences encoding mRNA, tRNA and rRNA are included within this definition. The sequences may be in the sense or antisense orientation with respect to the promoter. Antisense constructs can be used to
25 inhibit the expression of a gene in a cell according to well-known techniques. Nucleic acids may be, for example, ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or analogues thereof. Sequences encoding mRNA will optionally include some or all of 5' and/or 3' transcribed but untranslated flanking sequences naturally, or otherwise, associated with the translated coding sequence. It may optionally further include the
30 associated transcriptional control sequences normally associated with the transcribed sequences, for example transcriptional stop signals, polyadenylation sites and

downstream enhancer elements. Nucleic acids may comprise cDNA or genomic DNA (which may contain introns).

The basic structure of a retrovirus genome is a 5' LTR and a 3' LTR, between or
5 within which are located a packaging signal to enable the genome to be packaged, a
primer binding site, integration sites to enable integration into a host cell genome and
gag, *pol* and *env* genes encoding the packaging components - these are polypeptides
required for the assembly of viral particles. More complex retroviruses have
additional features, such as *rev* and RRE sequences in HIV, which enable the efficient
10 export of RNA transcripts of the integrated provirus from the nucleus to the cytoplasm
of an infected target cell.

In the provirus, these genes are flanked at both ends by regions called long terminal
repeats (LTRs). The LTRs are responsible for proviral integration, and transcription.
15 LTRs also serve as enhancer-promoter sequences and can control the expression of the
viral genes. Encapsidation of the retroviral RNAs occurs by virtue of a *psi* sequence
located at the 5' end of the viral genome.

The LTRs themselves are identical sequences that can be divided into three elements,
20 which are called U3, R and U5. U3 is derived from the sequence unique to the 3' end
of the RNA. R is derived from a sequence repeated at both ends of the RNA and U5 is
derived from the sequence unique to the 5' end of the RNA. The sizes of the three
elements can vary considerably among different retroviruses.

25 In a defective retroviral vector genome *gag*, *pol* and *env* may be absent or not
functional. The R regions at both ends of the RNA are repeated sequences. U5 and
U3 represent unique sequences at the 5' and 3' ends of the RNA genome respectively.

Preferred vectors for use in accordance with one aspect of the present invention are
30 recombinant non-primate lentiviral vectors.

The term "recombinant lentiviral vector" (RLV) refers to a vector with sufficient retroviral genetic information to allow packaging of an RNA genome, in the presence of packaging components, into a viral particle capable of infecting a target cell. Infection of the target cell includes reverse transcription and integration into the target cell genome. The RLV carries non-viral coding sequences which are to be delivered by the vector to the target cell. An RLV is incapable of independent replication to produce infectious retroviral particles within the final target cell. Usually the RLV lacks a functional *gag-pol* and/or *env* gene and/or other genes essential for replication. The vector of the present invention may be configured as a split-intron vector. A split intron vector is described in PCT patent application WO 99/15683.

Preferably the lentiviral vector of the present invention has a minimal viral genome.

As used herein, the term "minimal viral genome" means that the viral vector has been manipulated so as to remove the non-essential elements and to retain the essential elements in order to provide the required functionality to infect, transduce and deliver a nucleotide sequence of interest to a target host cell. Further details of this strategy can be found in our WO98/17815.

A minimal lentiviral genome for use in the present invention will therefore comprise (5') R - U5 - one or more first nucleotide sequences - U3-R (3'). However, the plasmid vector used to produce the lentiviral genome within a host cell/packaging cell will also include transcriptional regulatory control sequences operably linked to the lentiviral genome to direct transcription of the genome in a host cell/packaging cell. These regulatory sequences may be the natural sequences associated with the transcribed retroviral sequence, i.e. the 5' U3 region, or they may be a heterologous promoter such as another viral promoter, for example the CMV promoter. Some lentiviral genomes require additional sequences for efficient virus production. For example, in the case of HIV, *rev* and RRE sequence are preferably included. However the requirement for *rev* and RRE may be reduced or eliminated by codon optimisation. Further details of this strategy can be found in our WO01/79518.

In one embodiment of the present invention, the lentiviral vector is a self-inactivating vector.

By way of example, self-inactivating retroviral vectors have been constructed by deleting the transcriptional enhancers or the enhancers and promoter in the U3 region of the 3' LTR. After a round of vector reverse transcription and integration, these changes are copied into both the 5' and the 3' LTRs producing a transcriptionally inactive provirus (Yu *et al* 1986 Proc Natl Acad Sci 83: 3194-3198; Dougherty and Temin 1987 Proc Natl Acad Sci 84: 1197-1201; Hawley *et al* 1987 Proc Natl Acad Sci 84: 2406-2410; Yee *et al* 1987 Proc Natl Acad Sci 91: 9564-9568). However, any promoter(s) internal to the LTRs in such vectors will still be transcriptionally active. This strategy has been employed to eliminate effects of the enhancers and promoters in the viral LTRs on transcription from internally placed genes. Such effects include increased transcription (Jolly *et al* 1983 Nucleic Acids Res 11: 1855-1872) or suppression of transcription (Emerman and Temin 1984 Cell 39: 449-467). This strategy can also be used to eliminate downstream transcription from the 3' LTR into genomic DNA (Herman and Coffin 1987 Science 236: 845-848). This is of particular concern in human gene therapy where it is of critical importance to prevent the adventitious activation of an endogenous oncogene.

20

In our WO99/32646 we give details of features which may advantageously be applied to the present invention. In particular, it will be appreciated that the non-primate lentivirus genome (1) preferably comprises a deleted *gag* gene wherein the deletion in *gag* removes one or more nucleotides downstream of about nucleotide 350 or 354 of the *gag* coding sequence; (2) preferably has one or more accessory genes absent from the non-primate lentivirus genome; (3) preferably lacks the *tat* gene but includes the leader sequence between the end of the 5' LTR and the ATG of *gag*; and (4) combinations of (1), (2) and (3). In a particularly preferred embodiment the lentiviral vector comprises all of features (1) and (2) and (3).

30

The non-primate lentiviral vector may be a targeted vector. The term "targeted vector" refers to a vector whose ability to infect/transfect/transduce a cell or to be expressed in

a host and/or target cell is restricted to certain cell types within the host organism, usually cells having a common or similar phenotype.

5 Target cells for gene therapy using retroviral vectors include but are not limited to haematopoietic cells, (including monocytes, macrophages, lymphocytes, granulocytes, or progenitor cells of any of these); endothelial cells, tumour cells, stromal cells, astrocytes, or glial cells, muscle cells, epithelial cells, neurons, fibroblasts, hepatocyte, astrocyte, kidney, liver, heart and lung cells.

10 The vector may be pseudotyped with any molecule of choice, including but not limited to envelope glycoproteins (wild type or engineered variants or chimeras) of VSV-G, rabies, Mokola, MuLV, LCMV, Sendai, Ebola.

15 Although the first aspect of the present invention is directed to a method which, in particular, uses lentiviral vectors, other aspects of the present invention may employ other viral expression vectors. Viral vectors according to these aspects include but are not limited to a retroviral vector, a lentiviral vector, an adenoviral vector, an adeno-associated viral vector, a herpes viral vector, a pox viral vector, a parvoviral vector or a baculoviral vector.

20

The retroviral vector employed in the aspects of the present invention may be derived from or may be derivable from any suitable retrovirus. A large number of different retroviruses have been identified. Examples include: murine leukemia virus (MLV), human immunodeficiency virus (HIV), human T-cell leukemia virus (HTLV), mouse mammary tumour virus (MMTV), Rous sarcoma virus (RSV), Fujinami sarcoma virus (FuSV), Moloney murine leukemia virus (Mo-MLV), FBR murine osteosarcoma virus (FBR MSV), Moloney murine sarcoma virus (Mo-MSV), Abelson murine leukemia virus (A-MLV), Avian myelocytomatosis virus-29 (MC29), and Avian erythroblastosis virus (AEV). A detailed list of retroviruses may be found in Coffin *et al.*, 1997, "retroviruses", Cold Spring Harbour Laboratory Press Eds: JM Coffin, SM Hughes, HE Varmus pp 758-763.

25

30

Retroviruses may be broadly divided into two categories: namely, "simple" and "complex". Retroviruses may even be further divided into seven groups. Five of these groups represent retroviruses with oncogenic potential. The remaining two groups are the lentiviruses and the spumaviruses. A review of these retroviruses is presented in
5 Coffin *et al.*, 1997 (*ibid*).

The adenovirus is a double-stranded, linear DNA virus that does not go through an RNA intermediate. There are over 50 different human serotypes of adenovirus divided into 6 subgroups based on the genetic sequence homology. The natural target of
10 adenovirus is the respiratory and gastrointestinal epithelia, generally giving rise to only mild symptoms. Serotypes 2 and 5 (with 95% sequence homology) are most commonly used in adenoviral vector systems and are normally associated with upper respiratory tract infections in the young.

15 Viral gene expression can be divided into early (E) and late (L) phases. The late phase is defined by the onset of viral DNA replication. Adenovirus structural proteins are generally synthesised during the late phase. Following adenovirus infection, host cellular mRNA and protein synthesis is inhibited in cells infected with most serotypes. The adenovirus lytic cycle with adenovirus 2 and adenovirus 5 is very efficient and
20 results in approximately 10,000 virions per infected cell along with the synthesis of excess viral protein and DNA that is not incorporated into the virion. Early adenovirus transcription is a complicated sequence of interrelated biochemical events but it entails essentially the synthesis of viral RNAs prior to the onset of DNA replication.

25 The organisation of the adenovirus genome is similar in all of the adenovirus groups and specific functions are generally positioned at identical locations for each serotype studied. Early cytoplasmic messenger RNAs are complementary to four defined, noncontiguous regions on the viral DNA. These regions are designated E1-E4. The early transcripts have been classified into an array of intermediate early (E1a), delayed
30 early (E1b, E2a, E2b, E3 and E4), and intermediate regions.

The early genes are expressed about 6-8 hours after infection and are driven from 7

promoters in gene blocks E1-4.

Adenoviruses may be converted for use as vectors for gene transfer by deleting the E1 gene, which is important for the induction of the E2, E3 and E4 promoters. The E1-
5 replication defective virus may be propagated in a cell line that provides the E1 polypeptides in trans, such as the human embryonic kidney cell line 293. A therapeutic gene or genes can be inserted by recombination in place of the E1 gene. Expression of the gene is driven from either the E1 promoter or a heterologous promoter.

10

Even more attenuated adenoviral vectors have been developed by deleting some or all of the E4 open reading frames (ORFs). However, certain second generation vectors appear not to give longer-term gene expression, even though the DNA seems to be maintained. Thus, it appears that the function of one or more of the E4 ORFs may be
15 to enhance gene expression from at least certain viral promoters carried by the virus.

An alternative approach to making a more defective virus has been to "gut" the virus completely maintaining only the terminal repeats required for viral replication. The "gutted" or "gutless" viruses can be grown to high titres with a first generation helper
20 virus in the 293 cell line but it has been difficult to separate the "gutted" vector from the helper virus.

The adenovirus provides advantages as a vector for identifying candidate modulating moieties over other gene therapy vector systems for the following reasons:

25

It is a double stranded DNA nonenveloped virus that is capable of *in vivo* and *in vitro* transduction of a broad range of cell types of human and non-human origin. These cells include respiratory airway epithelial cells, hepatocytes, muscle cells, cardiac myocytes, synoviocytes, primary mammary epithelial cells and post-mitotically
30 terminally differentiated cells such as neurons.

Adenoviral vectors are also capable of transducing non dividing cells. This is very

important for diseases, such as cystic fibrosis, in which the affected cells in the lung epithelium, have a slow turnover rate. In fact, several trials are underway utilising adenovirus-mediated transfer of cystic fibrosis transporter (CFTR) into the lungs of afflicted adult cystic fibrosis patients.

5

The expression of viral or foreign genes from the adenovirus genome does not require a replicating cell. Adenoviral vectors enter cells by receptor mediated endocytosis. Once inside the cell, adenovirus vectors rarely integrate into the host chromosome. Instead, it functions episomally (independently from the host genome) as a linear
10 genome in the host nucleus. Hence the use of recombinant adenovirus alleviates the problems associated with random integration into the host genome.

Pox viral vectors may be used in accordance with aspects of the present invention, as large fragments of DNA are easily cloned into its genome and recombinant attenuated
15 vaccinia variants have been described (Meyer, *et al.*, 1991, J. Gen. Virol. 72: 1031-1038, Smith and Moss 1983 Gene, 25:21-28).

Examples of pox viral vectors include but are not limited to leporipoxvirus: Upton, *et al* J. Virology 60:920 (1986) (Shope fibroma virus); capripoxvirus: Gershon, *et al* J.
20 Gen. Virol. 70:525 (1989) (Kenya sheep-1); orthopoxvirus: Weir, *et al* J. Virol 46:530 (1983) (vaccinia); Esposito, *et al* Virology 135:561 (1984) (monkeypox and variola virus); Hruby, *et al* PNAS, 80:3411 (1983) (vaccinia); Kilpatrick, *et al* Virology 143:399 (1985) (Yaba monkey tumour virus); avipoxvirus: Binns, *et al* J. Gen. Virol 69:1275 (1988) (fowlpox); Boyle, *et al* Virology 156:355 (1987) (fowlpox);
25 Schnitzlein, *et al* J. Virological Method, 20:341 (1988) (fowlpox, quailpox); entomopox (Lytvyn, *et al* J. Gen. Virol 73:3235-3240 (1992)].

Poxvirus vectors are used extensively as expression vehicles for genes of interest in eukaryotic cells. Their ease of cloning and propagation in a variety of host cells has
30 led, in particular, to the widespread use of poxvirus vectors for expression of foreign protein and as delivery vehicles for vaccine antigens (Moss, B. 1991, Science 252: 1662-7).

Pox viruses which may be used in accordance with aspects of the present invention include but are not limited to recombinant pox viral vectors such as fowl pox virus (FPV), entomopox virus, vaccinia virus such as NYVAC, canarypox virus, MVA or
5 other non-replicating viral vector systems such as those described for example in WO 95/30018. Pox virus vectors have also been described where at least one immune evasion gene has been deleted (see WO 00/29428).

As indicated above, a nucleotide sequence used in a method of the present invention is
10 inserted into a vector which is operably linked to a control sequence that is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The NOI produced by a host recombinant cell may be secreted or may be contained intracellularly depending on the sequence and/or the vector used.

15 The heterologous gene, i.e. NOI, may be any allelic variant of a wild-type gene, or it may be a mutant gene. The term "gene" is intended to cover nucleic acid sequences which are capable of being at least transcribed. Thus, sequences encoding mRNA, tRNA and rRNA are included within this definition. The sequences may be in the sense or antisense orientation with respect to the promoter. Antisense constructs can be
20 used to inhibit the expression of a gene in a cell according to well-known techniques. Nucleic acids may be, for example, ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or analogues thereof. Sequences encoding mRNA will optionally include some or all of 5' and/or 3' transcribed but untranslated flanking sequences naturally, or otherwise, associated with the translated coding sequence. It may optionally further
25 include the associated transcriptional control sequences normally associated with the transcribed sequences, for example transcriptional stop signals, polyadenylation sites and downstream enhancer elements. Nucleic acids may comprise cDNA or genomic DNA (which may contain introns). However, it is generally preferred to use cDNA because it is expressed more efficiently since intron removal is not required.

30

Suitable NOI coding sequences include those that are of therapeutic and/or diagnostic application such as, but are not limited to: sequences encoding cytokines, chemokines,

hormones, antibodies, engineered immunoglobulin-like molecules, a single chain antibody, fusion proteins, enzymes, immune co-stimulatory molecules, immunomodulatory molecules, anti-sense RNA, a transdominant negative mutant of a target protein, a toxin, a conditional toxin, an antigen, a tumour suppressor protein and growth factors, membrane proteins, vasoactive proteins and peptides, anti-viral proteins and ribozymes, and derivatives thereof (such as with an associated reporter group). When included, such coding sequences may be typically operatively linked to a suitable promoter, which may be a promoter driving expression of a ribozyme(s), or a different promoter or promoters.

Suitable NOIs for use in the present invention in the treatment or prophylaxis of cancer include NOIs encoding proteins which: destroy the target cell (for example a ribosomal toxin), act as: tumour suppressors (such as wild-type p53); activators of anti-tumour immune mechanisms (such as cytokines, co-stimulatory molecules and immunoglobulins); inhibitors of angiogenesis; or which provide enhanced drug sensitivity (such as pro-drug activation enzymes); indirectly stimulate destruction of target cell by natural effector cells (for example, strong antigen to stimulate the immune system or convert a precursor substance to a toxic substance which destroys the target cell (for example a prodrug activating enzyme)). Encoded proteins could also destroy bystander tumour cells (for example with secreted antitumour antibody-ribosomal toxin fusion protein), indirectly stimulate destruction of bystander tumour cells (for example cytokines to stimulate the immune system or procoagulant proteins causing local vascular occlusion) or convert a precursor substance to a toxic substance which destroys bystander tumour cells (e.g. an enzyme which activates a prodrug to a diffusible drug).

NOI(s) may be used which encode antisense transcripts or ribozymes which interfere with expression of cellular genes for tumour persistence (for example against aberrant *myc* transcripts in Burkitts lymphoma or against *bcr-abl* transcripts in chronic myeloid leukemia). The use of combinations of such NOIs is also envisaged.

For further information on the nature of therapeutic genes see WO95/21927 and WO98/15294.

Suitable NOIs for use in the treatment or prevention of ischaemic heart disease include
5 NOIs encoding plasminogen activators. Suitable NOIs for the treatment or prevention
of rheumatoid arthritis or cerebral malaria include genes encoding anti-inflammatory
proteins, antibodies directed against tumour necrosis factor (TNF) alpha, and anti-
adhesion molecules (such as antibody molecules or receptors specific for adhesion
molecules).

10

Examples of hypoxia regulatable therapeutic NOIs can be found in WO95/21927.

The NOI coding sequence may encode a fusion protein or a segment of a coding
sequence.

15

Instead of, or as well as, being selectively expressed in target tissues, the NOI or NOIs
may encode a pro-drug activating enzyme or enzymes which have no significant effect
or no deleterious effect until the individual is treated with one or more pro-drugs upon
which the enzyme or enzymes act. In the presence of the active NOI, treatment of an
20 individual with the appropriate pro-drug leads to enhanced reduction in tumour growth
or survival.

A pro-drug activating enzyme may be delivered to a tumour site for the treatment of a
cancer. In each case, a suitable pro-drug is used in the treatment of the patient in
25 combination with the appropriate pro-drug activating enzyme. An appropriate pro-
drug is administered in conjunction with the vector. Examples of pro-drugs include:
etoposide phosphate (with alkaline phosphatase); 5-fluorocytosine (with cytosine
deaminase); doxorubicin-N-p-hydroxyphenoxyacetamide (with penicillin-V-amidase);
para-N-bis(2-chloroethyl) aminobenzoyl glutamate (with carboxypeptidase G2);
30 cephalosporin nitrogen mustard carbamates (with β -lactamase); SR4233 (with P450
Reductase); ganciclovir (with HSV thymidine kinase); mustard pro-drugs with
nitroreductase and cyclophosphamide (with P450).

Examples of suitable pro-drug activating enzymes for use in the invention include a thymidine phosphorylase which activates the 5-fluoro-uracil pro-drugs capcetabine and furtulon; thymidine kinase from herpes simplex virus which activates ganciclovir;
5 a cytochrome P450 which activates a pro-drug such as cyclophosphamide to a DNA damaging agent; and cytosine deaminase which activates 5-fluorocytosine. Preferably, an enzyme of human origin is used.

Suitable NOIs for use in the treatment or prevention of ischaemic heart disease include
10 NOIs encoding plasminogen activators. Suitable NOIs for the treatment or prevention of rheumatoid arthritis or cerebral malaria include genes encoding anti-inflammatory proteins, antibodies directed against tumour necrosis factor (TNF) alpha, and anti-adhesion molecules (such as antibody molecules or receptors specific for adhesion molecules).

15 The expression products encoded by the NOIs may be proteins which are secreted from the cell. Alternatively the NOI expression products are not secreted and are active within the cell. In either event, it is preferred for the NOI expression product to demonstrate a bystander effect or a distant bystander effect; that is the production of
20 the expression product in one cell leading to the killing of additional, related cells, either neighbouring or distant (e.g. metastatic), which possess a common phenotype.

Where macrophages or other haematopoietic cells are used, NOIs may be used which encode, for example, cytokines. These would serve to direct the subsequent
25 differentiation of the haematopoietic stem cells (HSCs) containing a therapeutic NOI. Suitable cytokines and growth factors include but are not limited to: ApoE, Apo-SAA, BDNF, Cardiotrophin-1, EGF, ENA-78, Eotaxin, Eotaxin-2, Exodus-2, FGF-acidic, FGF-basic, fibroblast growth factor-10, FLT3 ligand, Fractalkine (CX3C), GDNF, G-CSF, GM-CSF, GF- β 1, insulin, IFN- γ , IGF-I, IGF-II, IL-1 α , IL-1 β , IL-2, IL-3, IL-4,
30 IL-5, IL-6, IL-7, IL-8 (72 a.a.), IL-8 (77 a.a.), IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18 (IGIF), Inhibin α , Inhibin β , IP-10, keratinocyte growth factor-2 (KGF-2), KGF, Leptin, LIF, Lymphotactin, Mullerian inhibitory substance, monocyte

colony inhibitory factor, monocyte attractant protein, M-CSF, MDC (67 a.a.), MDC (69 a.a.), MCP-1 (MCAF), MCP-2, MCP-3, MCP-4, MDC (67 a.a.), MDC (69 a.a.), MIG, MIP-1 α , MIP-1 β , MIP-3 α , MIP-3 β , MIP-4, myeloid progenitor inhibitor factor-1 (MPIF-1), NAP-2, Neurturin, Nerve growth factor, β -NGF, NT-3, NT-4, Oncostatin M, PDGF-AA, PDGF-AB, PDGF-BB, PF-4, RANTES, SDF1 α , SDF1 β , SCF, SCGF, stem cell factor (SCF), TARC, TGF- α , TGF- β , TGF- β 2, TGF- β 3, tumour necrosis factor (TNF), TNF- α , TNF- β , TNIL-1, TPO, VEGF, GCP-2, GRO/MGSA, GRO- β , GRO- γ and HCC1.

For some applications, a combination of some of these cytokines may be preferred, in particular a combination which includes IL-3, IL-6 and SCF, for the maintenance and expansion of stem cell populations. For differentiation *in vitro*, further cytokines may be added such as GM-CSF and M-CSF to induce differentiation of macrophages or GM-CSF and G-CSF to obtain neutrophils. On reintroduction of the engineered cells into the individual from whom they were derived, the body's own mechanisms then permit the cells or their differentiated progeny to migrate into the affected area e.g. the tumour.

Optionally, another NOI may be a suicide gene, expression of which in the presence of an exogenous substance results in the destruction of the transfected or transduced cell. An example of a suicide gene includes the herpes simplex virus thymidine kinase gene (HSV *tk*) which can kill infected and bystander cells following treatment with ganciclovir.

Optionally another NOI may be a targeting protein (such as an antibody to the stem cell factor receptor (WO-A-92/17505; WO-A-92/21766)). For example, recombinant (ecotropic) retroviruses displaying an antibody (or growth factor or peptide) against a receptor present on HSCs (CD34 or stem cell factor, for example) might be used for targeted cell delivery to these cells, either *ex vivo* by incubating unfractionated bone marrow with virus or by intravenous delivery of virus.

NOIs may also include marker genes (for example encoding β -galactosidase or green fluorescent protein) or genes whose products regulate the expression of other genes. In addition, NOIs may comprise sequences coding fusion protein partners in frame with a sequence encoding a protein of interest. Examples of fusion protein partners
5 include the DNA binding or transcriptional activation domain of GAL4, a 6xHis tag and β -galactosidase. It may also be desirable to add targeting sequences to target proteins encoding by NOIs to particular cell compartments or to secretory pathways. Such targeting sequences have been extensively characterised in the art.

10 In one embodiment, at least one NOI, operably linked to a bacterial HRE according to the present invention encodes an oxygen-responsive bacterial transcriptional regulatory protein such as FNR. Such a construct will provide an autoregulated system since in the presence of hypoxia, expression of the bacterial transcriptional regulatory protein from the HRE construct will increase and serve to further increase
15 transcription from the HRE construct and other HRE constructs present.

In one preferred embodiment, the NOI encodes a ribozyme. Ribozymes are RNA molecules that can function to catalyse specific chemical reactions within cells without the obligatory participation of proteins. For example, group I ribozymes take the form
20 of introns which can mediate their own excision from self-splicing precursor RNA. Other ribozymes are derived from self-cleaving RNA structures which are essential for the replication of viral RNA molecules. Like protein enzymes, ribozymes can fold into secondary and tertiary structures that provide specific binding sites for substrates as well as cofactors, such as metal ions. Examples of such structures include
25 hammerhead, hairpin or stem-loop, pseudoknot and hepatitis delta antigenomic ribozymes have been described.

Each individual ribozyme has a motif which recognises and binds to a recognition site in a target RNA. This motif takes the form of one or more "binding arms" but
30 generally two binding arms. The binding arms in hammerhead ribozymes are the flanking sequences Helix I and Helix III which flank Helix II. These can be of variable length, usually between 6 to 10 nucleotides each, but can be shorter or longer.

The length of the flanking sequences can affect the rate of cleavage. For example, it has been found that reducing the total number of nucleotides in the flanking sequences from 20 to 12 can increase the turnover rate of the ribozyme cleaving a HIV sequence, by 10-fold (Goodchild, JVK, 1991 Arch Biochem Biophys 284: 386-391). A catalytic motif in the ribozyme Helix II in hammerhead ribozymes cleaves the target RNA at a site which is referred to as the cleavage site. Whether or not a ribozyme will cleave any given RNA is determined by the presence or absence of a recognition site for the ribozyme containing an appropriate cleavage site.

- Each type of ribozyme recognizes its own cleavage site. The hammerhead ribozyme cleavage site has the nucleotide base triplet GUX directly upstream where G is guanine, U is uracil and X is any nucleotide base. Hairpin ribozymes have a cleavage site of BCUGNYR, where B is any nucleotide base other than adenine, N is any nucleotide, Y is cytosine or thymine and R is guanine or adenine. Cleavage by hairpin ribozymes takes places between the G and the N in the cleavage site.

More details on ribozymes may be found in "Molecular Biology and Biotechnology" (Ed. RA Meyers 1995 VCH Publishers Inc p831-8320 and in "Retroviruses" (Ed. JM Coffin et al 1997 Cold Spring Harbour Laboratory Press pp 683).

20

Expression of the ribozyme may be induced in all cells, but will only exert an effect in those in which the target gene transcript is present.

- Alternatively, instead of preventing the association of the components directly, the substance may suppress the biologically available amount of a polypeptide of the invention. This may be by inhibiting expression of the component, for example at the level of transcription, transcript stability, translation or post-translational stability. An example of such a substance would be antisense RNA or double-stranded interfering RNA sequences which suppresses the amount of mRNA biosynthesis.

- In another preferred embodiment, the NOI comprises an siRNA. Post-transcriptional

gene silencing (PTGS) mediated by double-stranded RNA (dsRNA) is a conserved cellular defence mechanism for controlling the expression of foreign genes. It is thought that the random integration of elements such as transposons or viruses causes the expression of dsRNA which activates sequence-specific degradation of homologous single-stranded mRNA or viral genomic RNA. The silencing effect is known as RNA interference (RNAi). The mechanism of RNAi involves the processing of long dsRNAs into duplexes of 21-25 nucleotide (nt) RNAs. These products are called small interfering or silencing RNAs (siRNAs) which are the sequence-specific mediators of mRNA degradation. In differentiated mammalian cells dsRNA >30bp has been found to activate the interferon response leading to shut-down of protein synthesis and non-specific mRNA degradation (Stark et al 1998). However this response can be bypassed by using 21nt siRNA duplexes (Elbashir et al 2001, Hutvagner et al 2001) allowing gene function to be analysed in cultured mammalian cells.

15

In one embodiment an RNA polymerase III promoter, e.g., U6, whose activity is regulated by the presence of tetracycline may be used to regulate expression of the siRNA (Ohkawa et al, 2000).

20 In another embodiment the NOI comprises a micro-RNA. Micro-RNAs are a very large group of small RNAs produced naturally in organisms, at least some of which regulate the expression of target genes. Founding members of the micro-RNA family are *let-7* and *lin-4*. The *let-7* gene encodes a small, highly conserved RNA species that regulates the expression of endogenous protein-coding genes during worm development. The active RNA species is transcribed initially as an ~70nt precursor, which is post- transcriptionally processed into a mature ~21nt form. Both *let-7* and *lin-4* are transcribed as hairpin RNA precursors which are processed to their mature forms by Dicer enzyme (Lagos-Quintana et al, 2001).

25 30 In a further embodiment the NOI comprises double-stranded interfering RNA in the form of a hairpin. The short hairpin may be expressed from a single promoter, e.g., U6. In an alternative embodiment an effective RNAi may be mediated by incorporating

two promoters, e.g., U6 promoters, one expressing a region of sense and the other the reverse complement of the same sequence of the target. This is described in Example 9. In a further embodiment effective or double-stranded interfering RNA may be mediated by using two opposing promoters to transcribe the sense and antisense regions of the target from the forward and complementary strands of the expression cassette. These embodiments are described further in Example 9

In another embodiment the NOI may encode a short RNA which may act to redirect splicing ('exon-skipping') or polyadenylation or to inhibit translation. In the case of muscular dystrophy frame-shifting mutations in the dystrophin gene lead to a more severe Duchenne muscular dystrophy (DMD) phenotype than those which do not disrupt the translational reading frame (Becker muscular dystrophy). Antisense sequences targeted to induce skipping of exon 46 have been found to be effective in restoring dystrophin expression from the endogenous gene in DMD patient-derived muscle cells (van Deutekom et al 2001). Re-direction of polyadenylation by targeting antisense oligonucleotides to the 3' most polyadenylation site of E-selectin has been shown to re-direct polyadenylation to cryptic upstream sites resulting in transcripts with fewer instability sequences thereby increasing mRNA stability and altering protein expression (Vickers et al 2001). In this way the use of antisense can be applied to increase the abundance of a message. Targeting to sites crucial for initiation of translation is also possible, in this case the mRNA abundance is increased but protein levels decrease.

The NOI may be under the expression control of an expression regulatory element, usually a promoter or a promoter and enhancer. The enhancer and/or promoter may be preferentially active in a hypoxic or ischaemic or low glucose environment, such that the NOI is preferentially expressed in the particular tissues of interest, such as in the environment of a tumour, arthritic joint or other sites of ischaemia. Thus any significant biological effect or deleterious effect of the NOI on the individual being treated may be reduced or eliminated. The enhancer element or other elements conferring regulated expression may be present in multiple copies. Likewise, or in addition, the enhancer and/or promoter may be preferentially active in one or more

specific cell types - such as any one or more of macrophages, endothelial cells or combinations thereof. Further examples include include respiratory airway epithelial cells, hepatocytes, muscle cells, cardiac myocytes, synoviocytes, primary mammary epithelial cells and post-mitotically terminally differentiated non-replicating cells such as macrophages and neurons.

The term "operably linked" means that the components described are in a relationship permitting them to function in their intended manner. A library comprising a regulatory sequence "operably linked" to a reporter sequence is ligated in such a way that expression of the nucleic acid reporter sequence is achieved under conditions compatible with the control sequences.

The term "promoter" is used in the normal sense of the art, e.g. an RNA polymerase binding site in the Jacob-Monod theory of gene expression.

The term "enhancer" includes a DNA sequence which binds to other protein components of the transcription initiation complex and thus facilitates the initiation of transcription directed by its associated promoter.

The promoter and enhancer of the transcription units encoding the secondary delivery system are preferably strongly active, or capable of being strongly induced, in the primary target cells under conditions for production of the secondary delivery system. The promoter and/or enhancer may be constitutively efficient, or may be tissue or temporally restricted in their activity. Examples of temporally restricted promoters/enhancers are those which are responsive to ischaemia and/or hypoxia, such as hypoxia response elements or the promoter/enhancer of a *grp78* or a *grp94* gene. One preferred promoter-enhancer combination is a human cytomegalovirus (hCMV) major immediate early (MIE) promoter/enhancer combination.

In one preferred embodiment the combined use of a strong constitutive promoter such as CMV, or house-keeping promoter such as PGK, and the Tet-regulation system may

be used for control of gene expression. In addition to the Tet system other inducible systems include the metallothionein, hsp68, lacZ, and SV40 T antigen systems.

Transactivating factors may be employed through use of two transgenic lines, namely
5 one line which expresses the NOI under promoter "a", and a second line which expresses the transactivating factor "b" of promoter "a".

In another embodiment use may be made of the FLP recombinase system in which an inactive transgene is converted into the active form in a recombination event mediated
10 by yeast FLP recombinase. Use may also be made of the bacteriophage P1 Cre recombinase system, which allows genes to be silenced in particular cell or tissue types and at specific times of the organisms development.

Ubiquitous expression may be achieved using promoters from housekeeping genes,
15 such as beta-actin, mouse metallothionein, HMGR and histone H4.

Preferably the promoters of the present invention are tissue specific. That is, they are capable of driving transcription of an NOI in one tissue while remaining largely
"silent" in other tissue types.

20

The term "tissue specific" means a promoter which is not restricted in activity to a single tissue type but which nevertheless shows selectivity in that they may be active in one group of tissues and less active or silent in another group.

25 The level of expression of an NOI under the control of a particular promoter may be modulated by manipulating the promoter region. For example, different domains within a promoter region may possess different gene regulatory activities. The roles of these different regions are typically assessed using vector constructs having different variants of the promoter with specific regions deleted (that is, deletion analysis). This
30 approach may be used to identify, for example, the smallest region capable of conferring tissue specificity.

A number of tissue specific promoters, described above, may be particularly advantageous in practising the present invention. In most instances, these promoters may be isolated as convenient restriction digestion fragments suitable for cloning in a selected vector. Alternatively, promoter fragments may be isolated using the
5 polymerase chain reaction. Cloning of the amplified fragments may be facilitated by incorporating restriction sites at the 5' end of the primers.

Promoters suitable for cardiac-specific expression include the promoter from the murine cardiac α -myosin heavy chain (MHC) gene. Suitable vascular endothelium-
10 specific promoters include the Et-1 promoter and von Willebrand factor promoter.

Prostate specific promoters include the 5'flanking region of the human glandular kallikrein-1 (hKLK2) gene and the prostate specific antigen (hKLK3).

15 Examples of promoters/enhancers which are cell specific include a macrophage-specific promoter or enhancer, such as CSF-1 promoter-enhancer, or elements from a mannose receptor gene promoter-enhancer (Rouleux *et al* 1994 Exp Cell Res 214:113-119). Alternatively, promoter or enhancer elements which are preferentially active in neutrophils, or a lymphocyte-specific enhancer such as an IL-2 gene enhancer, may be
20 used.

Moreover, the NOI may be placed under the control of one or more sequences which confer developmentally-regulated expression. This will result in the NOIs being activated at a given stage in the development of the transgenic organism or its progeny.
25

Transcription of a NOI may be regulated by use of aptazymes. An aptazyme operably linked to a NOI may be activatable to cleave the transcript such that the NOI may be expressed following release of the NOI from the transcript. In a preferred embodiment the NOI is selected from the group comprising siRNA, short hairpin RNA, microRNA
30 and anti-sense RNA. For example, the addition of an aptazyme 5' of siRNA encoded by a short hairpin may allow the regulated induction (or inhibition) of self-cleavage of the transcript separating the hairpin from the aptazyme structure and hence activating

silencing. Ligands specific for the aptamer may be supplied exogenously, expressed endogenously or from the same vector. For example a protein ligand whose expression is controlled by the hypoxic response element (HRE) would only be synthesised under hypoxic conditions. If this were to activate cleavage of an aptazyme which released a
5 short hairpin targeted to vascular endothelial growth factor (VEGF), VEGF would be specifically down-regulated in hypoxia which would be therapeutically beneficial in a number of diseases including proliferative diabetic retinopathy. Alternatively the ligand for the aptamer could be VEGF itself.

10 In a further aspect of the invention, cleavage induced by the aptazyme may directly modulate expression of a NOI. The use of aptazymes in this way encompasses post-transcriptional regulation of a NOI according to the invention. The aptazyme may be activated (or inhibited) by the addition/removal of the appropriate ligand inducing cleavage of the transcript such that NOI expression is inhibited. The aptazyme may
15 cleave the transcript at for example the codon for the initiator methionine, or a UTR resulting in a transcript lacking either cap and/or poly-adenosine tail which will be subsequently degraded prior to translation.

This provides a means of shutting off synthesis of a NOI, for example a therapeutic
20 gene such as Factor IX, if levels are too high. Expression of the transgene may be engineered to be self-regulating in this way. For example an aptazyme whose activity is modulated by glucose binding could be designed such that high level expression of insulin occurs only when blood glucose levels are high. If glucose levels fall below a threshold level then the aptazyme is actived and the insulin transgene transcript
25 destroyed. An aptazyme which is regulated by doxycycline may be used to regulate the expression of NOIs both in vitro and in vivo by the administration of doxycycline.

The Tet-regulation system may be used to control expression of the aptazyme to provide an addition al level of control. Tet operator sequences inserted downstream of
30 the promoter may repress transcription in the presence of the Tet repressor protein

when doxycycline is removed, thereby preventing de novo transcription. As the aptazyme is active in the absence of doxycycline any existing transcripts will be cleaved and degraded.

- 5 The development of transgenic 'knockout' mouse technology has greatly benefited studies of gene function, with particular relevance in establishing mammalian models of genetic disease. Current technology is, however, limiting in certain cases. For example many genes, often those of medical significance, are essential for viability. In such cases pups die during embryonic development or soon after birth. The present invention provides an effective transgenic method for regulatable gene ablation such that the production of a protein of interest may be switched off at the desired developmental stage, facilitating the generation of disease models in adult mammals. The transgenic organism can then be out through one or more of any phenotype screen. Suitable general and directed phenotypic screens include the use of fundus photography, blood pressure, behaviour analysis, X-ray fluoroscopy, dual-energy X-ray absorptiometry (DEXA), CAT scans, complete blood counts (CBC), urinalysis, blood chemistry, insulin levels, glucose tolerance, fluorescence-activated cell sorting (FACS), histopathology, expression data, developmental biology. The methodology of the present invention will have broad application in many areas where temporal gene regulation would be advantageous and in validating putative drug targets identified in genomics programmes.

The present invention may be used to modulate the expression of genes that are associated with human disease. A non-exhaustive list of genes is set out below

- 25 (homologs of the genes are included):

Genes relating to cancer include, but are not limited to, *Cdh3*, *Ncam*, *Akp2*, *Asgr2*, *Bax*, *Bmp4*, *Ccnd1*, *Cd38*, *Cdc37*, *Cdkn1a*, *Cdkn1b*, *Cdkn1c*, *Csk*, *Epas1*, *Fgf2*, *Grpr*, *HBV*, *Igf1*, *Inhbb*, *Inpp5d*, *IRS1*, *Itga5*, *Kcna1*, *lacZ*, *Map2k4*, *Mdm2*, *Nfkb1a*, *Ngfb*, *Oxt*, *Pemt*, *Plp*, *Shh*, *Src*, *Stat5a*, *Tcfap2a*, *Trp53*, *Blmh*, *Cd152*, *Cmkar2*, *Cmkbr5*, *Csf1*, *Csf3*, *Egfr*, *Gzmb*, *Ifng*, *Ifngr*, *IGFBP3*, *Il1r1*, *Il1rap*, *Il2*, *Il2ra*, *Il2rb*, *Il2rg*, *Il4*, *Il4ra*, *Il5*, *Il6*, *Il7r*, *Il10*, *Il12a*, *Il12b*, *Il12rb1*, *Il12rb2*, *IRS1*, *Kdr*, *Lifr*, *Lta*, *Ncam*,

Ntf3, Ntf5, Ntrk1, Ntrk2, Ntrk3, Ph, Prlr, Scya3, Smst, Tgfa, Tgfb1, Tgfb2, Tgfb3, Tnf, Tnfrsf1a, Tnfrsf1b, Tnfrsf5, Apc, Prkdc, TAG, Amh, Kit, Kitl, Ter, Fech, hr, Atm, E2f1, Hox11, Apc, Cdh3, Erbb2, Hras, Met, Notch4, PIP, PyVT, Tag, Wnt1, Madh3, Nf1, Ptch, Rb1, Odc, Bcl3, Fos, Fyn, Jun, Kras2, luc, Mos, Myc, Rab3a, Rela, Yes, Cd44,
 5 *Mgmt, Plg, Ahr, Pgy2, Rag1, Btk, Igh-6, Jak3, Tcra, Tcrb, Tcrd, Tip53, Ttpa, Vhlh and Wt1.*

Genes relating to diabetes and obesity include, but are not limited to, *Ins2, Ins1, H2-Ea, H2-Ab1, Ifng, Prkdc, B2m, Rag1, Lep, Lepr, Cpe, Gck, Irs1, Irs2, Irs3, Irs4, Slc2a1, Cre, Dgat, tub, Pcsk2, Insr, Nos1, Nos3, Tnf, B2m, Thy1, Pomc, Ppara and*
 10 *Csf2.*

Genes relating to diseases of the cardiovascular system include, but are not limited to, *Acact, Alox15, Apoa2, Apob, Apoe, Ath1r, Cdkn1a, Cyp7a1, Epas1, Lcat, Ldlr, Pemt, Soat1, fld, hr, Ace, Adra1b, Adrb2, Adrbk1, Anx6, Atp7a, Cdh2, Evl, Fn1, Gjal, Itga4, Jup, Kif3a, Nf1, Nos3, Nppa, Thra, Vcam1, Wt1, Agt, Bdkrb2, Bmp4, Drd3,*
 15 *Kcna1, Npr3, Ren, Apoc1, Apoc2, Apoc3, Apoal, Cetp, Hpl, Lipc, Srb1, Adra2a, Agr1a, Fgf2, Tnf, Asgr2, Lrpap1, Vldlr, Col3a1 and Plg.*

Genes relating to diseases of the endocrine system include, but are not limited to, *A, Cpe, fld, Insr, Lep, Lepr, tub, Acact, acd, Cacnb4, Crh, Foxn1, gl, Bmp4, Csf1, dwg, fsn, Hcph, Kit, Kitl, Mitf, oc, Phex, Prlr, Sparc, Grpr, Amh, Ar, Cga, Fshb, jsd, Ghrhr,*
 20 *Hmgic, Myo5a, Nr5a1, Oxt, p, Pit1, Prop1, Smst, Agt, Igf1, Gck, Pcsk2, Egfr, Foxn1, Mclr, Tgfa, Thrbb, Tshr and Ttr.*

Genes relating to apoptosis include, but are not limited to, *Fas, Ngfr, Tnfrsf1a, Tnfrsf1b, Bax, Bcl2, E2f1, Mdm2, Pcc, Rb1, Trp53, Bdnf, Fas1, Gzmb, Ntf3, Ntf5, Pfp, Tag and Tnf.*

25 Genes relating to immunology and inflammation include, but are not limited to, *Cd1, Cd3e, Cd3z, Cd4, Cd44, Cd5, Cd8a, Cd8b, Cd14, Cd152, Cd28, Cd38, Fcgr1g, Fcgr2a, Fcgr2b, Fcgr3, Gp11, H2-Aa, H2-DMA, H2-Eb1, H2-Eb2, H2, Hc, Icam1, Igh-1, Igh-5, Igh, Igk-C, Igl-1, Igl-5, Itga4, Itga5, Itgb2, Itgp, Lyst, Mar1, Ncam, PCC, Pep3, Ptprc, Ptprcap, PVR, Sele, Sell, Selp, Spn, Taphh, Tcra, Tcrb, Tcrd, Thy1, Tlx1,*
 30 *Tnfrsf5, Tnfrsf6, Tnfsf5, Bmp4, Cmkar2, Cmkbr5, Csf1, Csf3, Egfr, Gzmb, Ifng, Ifngr,*

Il1r1, Il1rap, Il2, Il2ra, Il2rb, Il2rg, Il4, Il4ra, Il5, Il6, Il7r, Il10, Il12a, Il12b, Il12rb1, Il12rb2, Il15ra, Irs1, Itgb7, Kdr, Kitl, Lifr, Lta, Map2k4, Ntf3, Ntf5, Ntrk1, Ntrk3, Ph, Scya3, Smst, Tgfa, Tgfb1, Tgfb2, Tgfb3, Tnf, Tnfrsf1a, Tnfrsf1b, A, Atm, C3, C4, Cacnb4, Cd80, Cd86, Dh, Dsg3, Eef1a2, gl, hr, Lama2, Lbp, Lep, Lepr, Mitf, Pit1, Prop1, Scn8a, Abcb2, Ada, B2m, Bcl2, Bcl3, Btk, C2ta, Foxn1, H2-Ab1, Hcph, Igh-6, Igh-J, Ii, Jak3, Kit, Lck, Ltb, Lyn, Nfkb1, Nfkb1a, Pfp, Pnlliprp2, Prkdc, Ptprcap, Rag1, Relb, Stat4, Stat6, Tlr4, Alox5, Alox5ap, Alox15, Bdkrb2, Blmh, Bmp6, Cmo, Crh, Nos2, Ptg2, Vr1, Bax, E2f1, Inpp5d, Rb1, Stat5a, Trp53, Fyn and Irf1.

Genes relating to neurobiology include, but are not limited to, *Apoe, Atm, Bdnf, Cdk5, Chrna7, Cmkar4, Cstb, Gad2, Gfap, Gria2, Grik2, HD, Hdh, Nos1, Ntf3, Penk-rs, Prkcc, Psen1, Snca, Tnf and Vr1.*

In addition to the therapeutic gene or genes and the expression regulatory elements described, the delivery system may contain additional genetic elements for the efficient or regulated expression of the gene or genes, including promoters/enhancers, translation initiation signals, internal ribosome entry sites (IRES), splicing and polyadenylation signals. Expression levels may be improved by incorporating elements such as the WPRE.

The delivery of one or more one or more therapeutic genes by a delivery system according to the present invention may be used alone or in combination with other treatments or components of the treatment. In a further preferred embodiment of the first aspect of the invention, one or more nucleotides of interest (NOI) is introduced into the vector at the cloning site. Such therapeutic genes may be expressed from a promoter placed in the retroviral LTR or may be expressed from an internal promoter introduced at the cloning site.

For example, the delivery system of the present invention may be used to deliver one or more NOI(s) useful in the treatment of the disorders listed in WO98/05635. For ease of reference, part of that list is now provided: cancer, inflammation or inflammatory disease, dermatological disorders, fever, cardiovascular effects,

haemorrhage, coagulation and acute phase response, cachexia, anorexia, acute infection, HIV infection, shock states, graft-versus-host reactions, autoimmune disease, reperfusion injury, meningitis, migraine and aspirin-dependent anti-thrombosis; tumour growth, invasion and spread, angiogenesis, metastases, malignant,
5 ascites and malignant pleural effusion; cerebral ischaemia, ischaemic heart disease, osteoarthritis, rheumatoid arthritis, osteoporosis, asthma, multiple sclerosis, neurodegeneration, Alzheimer's disease, atherosclerosis, stroke, vasculitis, Crohn's disease and ulcerative colitis; periodontitis, gingivitis; psoriasis, atopic dermatitis, chronic ulcers, epidermolysis bullosa; corneal ulceration, retinopathy and surgical
10 wound healing; rhinitis, allergic conjunctivitis, eczema, anaphylaxis; restenosis, congestive heart failure, endometriosis, atherosclerosis or endosclerosis.

In addition, or in the alternative, the delivery system of the present invention may be used to deliver one or more NOI(s) useful in the treatment of disorders listed in
15 WO98/07859. For ease of reference, part of that list is now provided: cytokine and cell proliferation/differentiation activity; immunosuppressant or immunostimulant activity (e.g. for treating immune deficiency, including infection with human immune deficiency virus; regulation of lymphocyte growth; treating cancer and many autoimmune diseases, and to prevent transplant rejection or induce tumour immunity);
20 regulation of haematopoiesis, e.g. treatment of myeloid or lymphoid diseases; promoting growth of bone, cartilage, tendon, ligament and nerve tissue, e.g. for healing wounds, treatment of burns, ulcers and periodontal disease and neurodegeneration; inhibition or activation of follicle-stimulating hormone (modulation of fertility); chemotactic/chemokinetic activity (e.g. for mobilising
25 specific cell types to sites of injury or infection); haemostatic and thrombolytic activity (e.g. for treating haemophilia and stroke); antiinflammatory activity (for treating e.g. septic shock or Crohn's disease); as antimicrobials; modulators of e.g. metabolism or behaviour; as analgesics; treating specific deficiency disorders; in treatment of e.g. psoriasis, in human or veterinary medicine.

30

In addition, or in the alternative, the delivery system of the present invention may be used to deliver one or more NOI(s) useful in the treatment of disorders listed in

WO98/09985. For ease of reference, part of that list is now provided: macrophage inhibitory and/or T cell inhibitory activity and thus, anti-inflammatory activity; anti-immune activity, i.e. inhibitory effects against a cellular and/or humoral immune response, including a response not associated with inflammation; inhibit the ability of
5 macrophages and T cells to adhere to extracellular matrix components and fibronectin, as well as up-regulated fas receptor expression in T cells; inhibit unwanted immune reaction and inflammation including arthritis, including rheumatoid arthritis, inflammation associated with hypersensitivity, allergic reactions, asthma, systemic lupus erythematosus, collagen diseases and other autoimmune diseases, inflammation
10 associated with atherosclerosis, arteriosclerosis, atherosclerotic heart disease, reperfusion injury, cardiac arrest, myocardial infarction, vascular inflammatory disorders, respiratory distress syndrome or other cardiopulmonary diseases, inflammation associated with peptic ulcer, ulcerative colitis and other diseases of the gastrointestinal tract, hepatic fibrosis, liver cirrhosis or other hepatic diseases,
15 thyroiditis or other glandular diseases, glomerulonephritis or other renal and urologic diseases, otitis or other oto-rhino-laryngological diseases, dermatitis or other dermal diseases, periodontal diseases or other dental diseases, orchitis or epididymo-orchitis, infertility, orchidial trauma or other immune-related testicular diseases, placental dysfunction, placental insufficiency, habitual abortion, eclampsia, pre-eclampsia and
20 other immune and/or inflammatory-related gynaecological diseases, posterior uveitis, intermediate uveitis, anterior uveitis, conjunctivitis, chorioretinitis, uveoretinitis, optic neuritis, intraocular inflammation, e.g. retinitis or cystoid macular oedema, sympathetic ophthalmia, scleritis, retinitis pigmentosa, immune and inflammatory components of degenerative fundus disease, inflammatory components of ocular
25 trauma, ocular inflammation caused by infection, proliferative vitreo-retinopathies, acute ischaemic optic neuropathy, excessive scarring, e.g. following glaucoma filtration operation, immune and/or inflammation reaction against ocular implants and other immune and inflammatory-related ophthalmic diseases, inflammation associated with autoimmune diseases or conditions or disorders where, both in the central nervous
30 system (CNS) or in any other organ, immune and/or inflammation suppression would be beneficial, Parkinson's disease, complication and/or side effects from treatment of Parkinson's disease, AIDS-related dementia complex HIV-related encephalopathy,

Devic's disease, Sydenham chorea, Alzheimer's disease and other degenerative diseases, conditions or disorders of the CNS, inflammatory components of stokes, post-polio syndrome, immune and inflammatory components of psychiatric disorders, myelitis, encephalitis, subacute sclerosing pan-encephalitis, encephalomyelitis, acute
5 neuropathy, subacute neuropathy, chronic neuropathy, Guillain-Barre syndrome, Sydenham chora, myasthenia gravis, pseudo-tumour cerebri, Down's Syndrome, Huntington's disease, amyotrophic lateral sclerosis, inflammatory components of CNS compression or CNS trauma or infections of the CNS, inflammatory components of muscular atrophies and dystrophies, and immune and inflammatory related diseases,
10 conditions or disorders of the central and peripheral nervous systems, post-traumatic inflammation, septic shock, infectious diseases, inflammatory complications or side effects of surgery, bone marrow transplanTation or other transplantation complications and/or side effects, inflammatory and/or immune complications and side effects of gene therapy, e.g. due to infection with a viral carrier, or inflammation associated with
15 AIDS, to suppress or inhibit a humoral and/or cellular immune response, to treat or ameliorate monocyte or leukocyte proliferative diseases, e.g. leukaemia, by reducing the amount of monocytes or lymphocytes, for the prevention and/or treatment of graft rejection in cases of transplantation of natural or artificial cells, tissue and organs such as cornea, bone marrow, organs, lenses, pacemakers, natural or artificial skin tissue.

20

The subject treated by the method of the present invention may be an animal subject. Preferably the subject is a mammalian subject, more preferably a human subject.

The present invention also provides a pharmaceutical composition for treating an
25 individual by gene therapy, wherein the composition comprises a therapeutically effective amount of the delivery system of the present invention and optionally comprising one or more deliverable therapeutic and/or diagnostic NOI(s). Since the delivery system is a viral delivery system then the composition may in addition or in the alternative comprise a viral particle produced by or obtained from same. The
30 pharmaceutical composition may be for human or animal usage. Typically, a physician will determine the actual dosage which will be most suitable for an individual subject and it will vary with the age, weight and response of the particular individual.

The composition may optionally comprise a pharmaceutically acceptable carrier, diluent, excipient or adjuvant. The choice of pharmaceutical carrier, excipient or diluent can be selected with regard to the intended route of administration and standard pharmaceutical practice. The pharmaceutical compositions may comprise as - or in addition to - the carrier, excipient or diluent any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilising agent(s), and other carrier agents that may aid or increase the viral entry into the target site (such as for example a lipid delivery system).

10

Where appropriate, the pharmaceutical compositions can be administered by any one or more of: inhalation, in the form of a suppository or pessary, topically in the form of a lotion, solution, cream, ointment or dusting powder, by use of a skin patch, orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents, or they can be injected parenterally, for example intracavernosally, intravenously, intramuscularly or subcutaneously. For parenteral administration, the compositions may be best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood. For buccal or sublingual administration the compositions may be administered in the form of tablets or lozenges which can be formulated in a conventional manner.

20

The delivery of one or more therapeutic genes by a delivery system according to the invention may be used alone or in combination with other treatments or components of the treatment.

25

The non-primate lentiviral vector particles of the present invention are typically generated in a suitable producer cell. Producer cells are generally mammalian cells but can be for example insect cells. A producer cell may be a packaging cell containing the virus structural genes, normally integrated into its genome. The packaging cell is then transfected with a nucleic acid encoding the vector genome, for

30

the production of infective, replication defective vector particles. Alternatively the producer cell may be co-transfected with nucleic acid sequences encoding the vector genome and the structural components, and/or with the nucleic acid sequences present on one or more expression vectors such as plasmids, adenovirus vectors, herpes viral
5 vectors or any method known to deliver functional DNA into target cells.

The vectors of the invention, for example, the lentiviral vectors of the first aspect of the invention, may be used to deliver an NOI to any prenatal cell. The term "prenatal" means occurring or present before birth. In one embodiment the method is applied to a
10 cell at the embryonic stage. The term embryo includes animals in the early stages of development up to birth (or hatching). As used herein the term "embryo" includes "pre-embryo", i.e. the structure formed after fertilisation of an ovum but before differentiation of embryonic tissue, and includes a zygote and blastocyte. The term also includes a fetal cell, i.e. an embryonic cell which is in the latter stages of
15 development. The present invention also encompasses delivery to a perinatal cell. The term "perinatal" refers to the period from about 3 months before to about one month after birth, and includes the neonatal period. The term "neonate" refers to the first few weeks following birth.

20 Generally vectors of the invention, for example, the lentiviral vectors of the first aspect of the invention may be used to deliver an NOI to any germ cell, including a primordial germ cell, or cell which is capable of giving rise to a germ line change. The term "germ cell" is the collective term for cells in the reproductive organ of multicellular organisms that divide by meiosis to produce gametes. The term
25 "gametes" refers to the haploid reproductive cells - in effect the ovum and sperm. However, as indicated above the present invention is also applicable to cells involved in gametogenesis and cells from structures in which gametogenesis take place, such as the ovary.

30 Gametogenesis will now be described in relation to mammals by way of example only. Vectors such as the lentiviral vector may be used to deliver an NOI to any of the cells of structures mentioned below. It will be appreciated that the equivalent processes in

non-mammalian organisms are also included in the present invention. In brief, gametogenesis is the process of forming gametes (by definition haploid, n) from diploid cells of the germ line. Spermatogenesis is the process of forming sperm cells by meiosis (in animals, by mitosis in plants) in specialized organs known as gonads (in males these are termed testes). After division the cells undergo differentiation to become sperm cells. Oogenesis is the process of forming an ovum (egg) by meiosis (in animals, by mitosis in the gametophyte in plants) in specialized gonads known as ovaries.

10 In spermatogenesis the sperms are formed from the male germ cells, spermatogonia, which line the inner wall of the seminiferous tubules in the testis. A single spermatogonium divides by mitosis to form the primary spermatocyte, each of which undergoes the initial division of meiosis to form two secondary spermatocytes. Each of these then undergoes a second meiotic division to form two spermatids, which mature into spermatozoa. The testis is composed of numerous seminiferous tubules, in whose walls spermatogenesis takes place. The primordial germ cells are formed in the germinal epithelium lining towards the outside of the tubule, and as cell divisions proceed the daughter cells move towards the lumen of the tubule. All these cells are nourished and supported by neighbouring Sertoli cells.

20

In oogenesis a primary oocyte is formed by differentiation of an oogonium and then undergoes the first division of meiosis to form a polar body and a secondary oocyte. Following fertilisation of the egg, the secondary oocyte undergoes the second meiotic division to form the mature ovum and a second polar body. The ovary contains many follicles composed of a developing egg surrounded by an outer layer of follicle cells. After ovulation the egg moves down the oviduct to the uterus.

It will be appreciated that the vector may be administered at one locality, but the NOI is expressed or its effects felt, in another cell of the organism, i.e. the site of administration may be different from the target cell. Cells into which the non-primate lentiviral vector may be administered include the examples of target cells listed above.

More preferably, the cell is at the embryonic stage, and for example is in utero, the lentiviral vector may be administered via the umbilical cord, placenta, or amniotic fluid, or by the intraperitoneal or intrahepatic routes. The introduction of the lentiviral vector is aided by the use of ultrasound.

5

The production of transgenic animals, using ES cells and otherwise, is well known in the art, and described for example in *Manipulating the Mouse Embryo*, 2nd Ed., by B. Hogan, R. Beddington, F. Costantini, and E. Lacy. Cold Spring Harbor Laboratory Press, 1994; *Transgenic Animal Technology*, edited by C. Pinkert. Academic Press, Inc., 1994; 10 *Gene Targeting: A Practical Approach*, edited by A. L. Joyner. Oxford University Press, 1995; *Strategies in Transgenic Animal Science*, edited by G. M. Monastersky and J. M. Robl. ASM Press, 1995; and *Mouse Genetics: Concepts and Applications*, by Lee M. Silver, Oxford University Press, 1995. A useful general textbook on this subject is Houdebine, *Transgenic animals – Generation and Use* (Harwood Academic, 1997) – 15 an extensive review of the techniques used to generate transgenic animals from fish to mice and cows.

Thus, for example, the present invention permits the introduction of heterologous DNA into, for example, fertilised mammalian ova by lentiviral infection. In one 20 embodiment the fertilised egg is collected from a donor mother at the one cell stage and the transduced cell is transferred to a foster mother. Integration which occurs at the one cell stage produces an organism which is a true transgenic, i.e. transgenic throughout, including the germ cells. If integration occurs at a later stage mosaics are produced. In a highly preferred method, developing embryos are infected with a 25 lentivirus containing the desired DNA, and transgenic animals produced from the infected embryo. Traditional transgenic methods have required that the embryonic cells are transformed ex vivo then reimplanted into the uterus. A significant advantage associated with the present invention is that the NOI can be introduced in utero. Another method which may be used to produce a transgenic animal involves 30 introducing a nucleic acid into pro-nuclear stage eggs by lentiviral infection. Injected eggs are then cultured before transfer into the oviducts of pseudopregnant recipients.

By way of a specific example for the construction of transgenic mammals, such as cows, nucleotide constructs comprising a sequence encoding a therapeutic protein are introduced using the method of the present invention into oocytes which are obtained from ovaries freshly removed from the mammal. The oocytes are aspirated from the follicles and allowed to settle before fertilisation with thawed frozen sperm capacitated with heparin and prefractionated by Percoll gradient to isolate the motile fraction.

The fertilised oocytes are centrifuged, for example, for eight minutes at 15,000 g to visualise the pronuclei for injection and then cultured from the zygote to morula or blastocyst stage in oviduct tissue-conditioned medium. This medium is prepared by using luminal tissues scraped from oviducts and diluted in culture medium. The zygotes must be placed in the culture medium within two hours following microinjection.

Oestrous is then synchronized in the intended recipient mammals, such as cattle, by administering coprostanol. Oestrous is produced within two days and the embryos are transferred to the recipients 5-7 days after estrous. Successful transfer can be evaluated in the offspring by Southern blot.

Alternatively, the desired constructs can be introduced into embryonic stem cells (ES cells) and the cells cultured to ensure modification by the transgene. The modified cells are then injected into the blastula embryonic stage and the blastulas replaced into pseudopregnant hosts. The resulting offspring are chimeric with respect to the ES and host cells, and nonchimeric strains which exclusively comprise the ES progeny can be obtained using conventional cross-breeding. This technique is described, for example, in WO91/10741.

Analysis of animals which may contain transgenic sequences would typically be performed by either PCR or Southern blot analysis following standard methods. If desired, the organism can be bred to homozygosity.

The use of the present invention to produce transgenic organism for use in gene therapy and in the production of disease models has been mentioned above. In particular, disease models allow experimental investigation of gene function. In
5 general transgenic organisms expressing novel genes or genes with a heterologous promoter represent gain-of-function mutations. Loss-of-function mutations can be created by gene targeting to create so-called "knockout" organisms. Transgenic organisms are also useful for the investigation of control regions and expression patterns. Transgenic organisms can also be used to identify novel genes using
10 techniques such as insertional mutation, gene traps and promoter traps. Transgenic animals also have agricultural applications, for example to bring genetic improvements to milk yield, body mass, milk composition, disease resistance etc. Transgenic animals are also useful in so-called pharmaceutical farming in which transgenic livestock are used as bioreactors for the production of therapeutic proteins.

15

By way of example, the regulated ablation of SMN (homozygous deletion of which results in pre-natal mortality) would provide a useful model of spinal muscular atrophy for gene therapy studies. A CFTR deficiency model is also a valuable application. Other putative candidates include: presenilin-1, RAR α , BDNF, VEGF and EGFR.

20

The analysis of resultant phenotypes can be carried out using standard techniques such as histological tissue analysis and microarray gene expression profiling.

In accordance with a preferred feature of the present invention a lentiviral vector, and
25 preferably an EIAV vector, is used to produce transgenic chickens that produce therapeutic or diagnostic proteins in their eggs.

The use of lentiviral vectors to produce transgenic avians allows the expression of genes throughout significant numbers of generations without the foreign gene
30 silencing observed with some retroviral vectors. For example Mizuarai *et al.* (2001) (Biochemical and Biophysics Research Communications; 286: 456-463) observed that

LTR driven expression from MLV based vectors in transgenic quail could not be observed, whilst expression from an internal Rous sarcoma virus (RSV) promoter was present in G₀, G₁ and G₂ birds.

- 5 A lentiviral vector encoding for a therapeutic protein or a protein of diagnostic use, for example an antibody or a fragment of an antibody, may be used to produce transgenic chickens. The antibody may be engineered to contain domains derived from more than one animal species or to contain domains that bind to different target molecules. The nucleotide coding sequence of the target gene can be altered to increase RNA
10 stability or RNA transcription levels without altering the amino acid sequence of the resultant protein.

Expression of the therapeutic/diagnostic gene may be from a constitutive promoter or from a promoter that confers tissue specific expression. For instance expression of the
15 target protein may be restricted to the reproductive structures (including the oviduct or reproductive tract) in such a way as to result in the target protein being present in eggs. Promoters or elements from promoters of genes for proteins found in egg white such as the ovalbumin, lysozyme, conalbumin and ovomucoid may be used. The expression of these genes is regulated by the steroid hormones but there is evidence for the
20 ovalbumin and conalbumin promoters that other cell specific transcription factors are also involved (Dierich *et al* EMBO J. 1987 Aug;6(8):2305-12). The ovalbumin gene promoter has been shown to have tissue specific silencing elements between -3200 and -2800 bp from the transcription site (Muramatsu *et al*. Mol Cell Biochem 1998 Aug;185(1-2):27-32), whereas a silencing element is present -2400bp from the
25 transcription site of the lysozyme gene (Bonifer *et al*. J Biol Chem. 1997 Oct 17;272(42):26075-8. Review). However Dierich *et al*. (1987) obtained some degree of cell specificity in a truncated ovalbumin extending from -1348 to -1. Some degree of steroid regulation was observed for a truncated ovalbumin extending from -425 to -1 in primary cultured chicken oviduct tubular gland cells (Dierich *et al*. 1987).

30

Lentiviral vectors encoding for therapeutic/diagnostic proteins are used to transduce cells in the blastoderm stage embryo in new-laid eggs by injection. Alternatively,

lentiviral vectors can be used to transduce earlier stage embryos using techniques such as those described in WO 90/13626 or similar published techniques to allow the embryo to develop normally.

- 5 In brief a uterine embryo is abstracted from a hen either manually or by inducing premature oviposition. The embryo is transduced with the lentiviral vector and then cultured to fruition. This allows cells of the embryo to be transduced whilst the number of cells present is relatively low and increases the number of birds produced in which the introduced gene is present in the germ line and is inherited.

10

The present invention also relates to the use of RNAi to enhance protein yield from transgenic avians

- 15 In order to maximise the expression of a desired protein in eggs, as described in example 8 below, RNAi may be used to decrease the proportion of abundant proteins normally present in the egg white such as ovalbumin, lysozyme, conalbumin and ovomucoid. Down-regulating the production of these proteins may result in a concomitant increase in the proportion of the desired protein in the egg giving improved yields.

20

This may be achieved by expressing siRNAs, or short hairpin pre-cursors, targeting the required mRNA from an EIAV vector. The vector may comprise siRNAs targeting one or more mRNAs at one or more sites within the mRNA.

- 25 Efficient targeting of proteins such as ovalbumin may result in eggs which are non-viable for breeding purposes. Transgenic roosters may therefore be identified and crossed to hens transgenic for a vector containing the transgene of interest in order to obtain female offspring with both traits which would be used as bioreactors for protein production. Regulation of the siRNAs (as described in example 10) would be an
30 alternative to this: silencing may be induced in laying hens when eggs are required for protein production, but not for breeding. This would also overcome any possible

deleterious effects of down-regulating expression of the target genes within the whole organism.

The present invention will now be described by way of further example with reference
5 to the following non-limiting Examples:

Example 1 - EIAV transduction of perinatal animals

Preparation of Vector

Vector was prepared by transient co-transfection of 293T human embryonic kidney
10 cells as previously described (Mitrophanous et al 1999). The EIAV vector genome, SMART2Z, expresses the β -galactosidase reporter gene from an internal CMV promoter. It contains the EIAV central polypurine tract (cPPT) (Stetor et al 1999) and the Woodchuck Hepatitis Post-Transcriptional Regulatory Element (WPRE) (Donello et al 1998).

15

Administration of vector

Vector was administered by injecting foetuses intra-vascularly as follows:
Under isoflurane anaesthesia a full depth midline laparotomy was performed to expose
the uteri of pregnant mice at 16 days gestation. For each foetus 2×10^7 T.U.
20 (transforming units) of vector was administered in a total volume of 20 μ l, using a 34-gauge needle (Hamilton), into a peripheral yolk sac vessel. Up to five foetuses were injected per dam. The laparotomy was closed by suturing layer to layer and mice allowed to recover in a warm cage.

25 Detection of gene transfer

Pups were born at 18-19 days after conception. Mice were sacrificed at various stages of development (3, 7, 14, 28 and 79 days post-injection) by euthanising with isoflurane anaesthesia and samples prepared for histology. Organs were placed in 100% ethanol solution for 2h prior to staining with X-gal solution (1mg/ml 5-bromo-

4-chloro-3-indolyl- β -D-galactopyranoside dissolved in dimethylsulphoxide, 5mM potassium ferricyanide, 5mM potassium ferrocyanide, 2mM magnesium chloride) for the β -galactosidase marker gene expressed by the vector. Stained tissue was fixed in 10% formaldehyde for 2h and paraffin embedded, sectioned and counterstained with neutral red. Staining showed transduction of a number of organs including liver, lung, heart, muscle, kidney, skeletal muscle and brain. The results are shown in Figures 1 to 11.

Expression levels did not decrease over the period of the study and clonal expansion of transduced cells was observed.

In addition to injection into the yolk sac vessel or umbilical vein, injection directly into the circulation, CSF or other tissue may be carried out, or into the amniotic fluid. The latter may be particularly appropriate when transduction of lung or skin tissue is desired.

Example 2 – Haemophilia

This Example is carried out following the methodology of Example 1. Haemophilia is a blood condition in which an essential clotting factor is either partly or completely missing. It is an X-linked recessive disorder. There are two types of haemophilia, the most common being haemophilia A, in which Factor VIII is lacking. In haemophilia B, Factor IX is lacking. EIAV is used to deliver factor VIII or IX by EIAV to the umbilical vein of haemophiliac foetus or hepatic portal vein of perinates.

Preparation of the vector

A vector such as those described in our co-pending GB0202403.1. In more detail:

pONY 8.4 series of vectors has a number of modifications which enable it to function as part of a transient or stable vector system totally independent of accessory proteins, with no detrimental effect on titre. Conventionally lentiviral vector genomes have required the presence of the viral protein rev in producer cells (transient or stable) in

order to obtain adequate titres. This includes current HIV vector systems as well as earlier EIAV vectors.

There are 3 modifications when compared with the pONY 8.1 series of vector genomes, these are:

5

- a) All the ATG motifs which are derived from gag and form part of the packaging signal have been modified to read ATTG. This allows the insertion of an open reading frame which can be driven by a promoter in the LTR.
- 10 b) The length of the genome i.e. distance between the R regions is closer to that seen in the wt virus (7.9kb).
- c) The 3' U3 region has been modified to include sequences from the moloney leukemia virus (MLV) U3 region, so upon transduction it can drive second open reading frame (ORF) in addition to the internal cassette,
- 15 In this example we have MLV but this could be any promoter.

Further details on modifying LTRs can be found in our WO96/37623 and WO98/17816.

- 20 Figure 12 is a schematic representation of EIAV genomes. These may be used for transfection in accordance with the method of the present invention. Upon transfection the 3' LTR will be copied to the 5' LTR. Figure 13 gives the total plasmid sequence of pONY8.1G. Figure 14 gives the total plasmid sequence of pONY8.4ZCG. Figure 15 gives the total plasmid sequence of pONY8.4GCZ. Figure 16 is a schematic representation of the hybrid U3 region. Figure 17 gives the sequence of the hybrid LTR.
- 25

The construction of EIAV/MLV Hybrid LTR Vectors

PCR was carried out as follows:

30

Product A = primers KM001 + KM003, with the pONY8.1Z as target.

Product B = primers KM004 + KM005, with the pHIT111 as target.

5 Product C = primers KM006 + KM002, with the pONY8.1Z as target.

The PCR products (A, B and C) were gel purified. A PCR reaction was set up using Product A and B (with primers KM001 and KM005) to give Product D. A PCR reaction was set up using Product B and C (with primers KM004 and KM002) to give
10 Product E. Product D and E were gel purified and used in a PCR reaction, as targets with primers KM001 and KM002 to give Product F. The PCR Product F was gel purified (approximately 1 kb). This was then cut with Sap I and subcloned into pONY8.1Z cut with Sap I. This gave the vector pONY8.1Zhyb shown in Figures 18 and 19. The 3' LTR of ELAV has now been replaced with an ELAV/MLV hybrid
15 LTR. The ELAV U3 has been almost replaced with the MLV U3 region. The ELAV 5' U3 sequences of the 3' LTR have been retained as these comprise the *att* site, that is the sequences needed for integration.

The primer sequences are shown below:

20

ELAV/MLV hybrid U3

KM001

CAAAGCATGCCTGCAGGAATTCG

25

KM002

GAGCGCAGCGAGTCAGTGAGCGAG

KM003

30 GCCAAACCTACAGGTGGGGTC

TTTCATTATAAAAACCCCTCATAAAAACCCACAG

KM004

CTGTGGGGTTTTTATGAGGGGTTTTATAATGAAAGACCCACCTGTAGGTT

5 TGGC

KM005

GAAGGGACTCAGACCGCAGAATCTGAGTGCCCCCGAGTGAGGGGTTGTG
GGCTCT

10

KM006

AGAGCCCACAACCCCTCACTCGGGGG
GCACTCAGATTCTGCGGTCTGAGTCCCTTC

15 Sequence of final PCR product.

EIAV PPT/U3

CAAAGCATGCCTGCAGGAATTCGATATCAAGCTTATCGATACCGTCGAATT
GGAAGAGCTTTAAATCCTGGCACATCTCATGTATCAATGCCTCAGTATGTT

20 TAGAAAAACAAGGGGGGAAGTGTGGGGTTTTTATGAGGGGTTTTATAA

MLV U3

TGAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCATTTT
GCAAGGCATGGAAAAATACATAACTGAGAATAGAGAAGTTCAGATCAAGG

25 TCAGGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGGT

AAGCAGTTCCTGCCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAAT

ATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCTGCCCCGGCTCAGGG

CCAAGAACAGATGGTCCCAGATGCGGTCCAGCCCTCAGCAGTTTCTAGA

GAACCATCAGATGTTTCCAGGGTGCCCCAAGGACCTGAAATGACCCTGTGC

30 CTTATTTGAACTAACCAATCAGTTCGCTTCTCGCTTCTGTTTCGCGCGCTTCT

GCTCCCCGAGCTCAATAAAAGAGCCCACAACCCCTCACTCGGG

MLV U3/EIAV R/U5

GGGCACTCAGATTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGAAAAGGC
 CTTTGTAATAAATAATTTCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTT
 CGAGATCCTACAGAGCTCATGCCTTGGCGTAATCATGGTCATAGCTGTTTC
 5 CTGTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAA
 GCATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTA
 ATTGCGTTGCGCTCACTGCCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAG
 CTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGCGTATTGG
 GCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTC

10

Haemophilia A

An EIAV viral vector (such as those described above) expressing Factor VIII (the wild type full length open reading frame (ORF) or a truncated B domain deleted wild type ORF or a codon optimised ORF or a truncated B domain deleted codon optimised
 15 ORF) is administered following the methodology in Example 1 and including either intravenous or intra hepatic or intramuscular delivery. A suitable promoter such as CMV or human promoter/enhancers such as PGK is used to express the gene. In addition inducible promoters such as the Tet system can be used to regulate the expression. As an alternative tissue specific promoter/enhancers can be used to limit
 20 expression to the cell types.

Haemophilia B

An EIAV viral vector (such as those described above) expressing Factor IX (the wild type ORF or a codon optimised ORF) is administered following the methodology in
 25 Example 1 and including either intravenous or intra hepatic or intramuscular delivery. A suitable promoter such as CMV or human promoter/enhancers such as PGK is used to express the gene. In addition inducible promoters such as the Tet system can be used to regulate the expression. As an alternative tissue specific promoter/enhancers can be used to limit expression to the cell types.

30

Example 3 - Cystic fibrosis

This Example is carried out following the methodology of Example 1. Cystic fibrosis is an hereditary recessive disorder caused by mutation of cystic fibrosis transmembrane conductance regulator (CFTR), a protein that is thought to have a role in ion transport, mucus rheology, inflammation and bacterial adherence. EIAV is used to deliver CFTR by to the amniotic fluid for transduction of lung.

An EIAV viral vector (such as those described above) expressing CFTR (the wild type ORF or a codon optimised ORF) is administered following the methodology in Example 1 and including either intragastrintestinal delivery intralung or intraamniotic fluid. A suitable promoter such as CMV or human promoter/enhancers such as PGK is used to express the gene. In addition inducible promoters such as the Tet system can be used to regulate the expression. As an alternative tissue specific promoter/enhancers can be used to limit expression to the cell types.

Example 4 - Muscular dystrophy

This Example is carried out following the methodology of Example 1. Duchenne muscular dystrophy (DMD) is a lethal X-linked recessive disorder. DMD results from genetic deficiency in the level and/or activity of the protein dystrophin in the striated musculature. EIAV is used to deliver of minidystrophin cDNA (corresponding to a mild Becker muscular dystrophy (BMD) phenotype) to the umbilical vein of perinates and/or directly into foetal skeletal muscle.

25

An EIAV viral vector (such as those described above) expressing dystrophin (the wild type full length open reading frame (ORF) or a truncated wild type ORF or a codon optimised ORF or a truncated codon optimised ORF) is administered following the methodology in Example 1 and including delivery into all muscle groups. A suitable promoter such as CMV or human promoter/enhancers such as PGK is used to express the gene. In addition inducible promoters such as the Tet system can be used to

30

regulate the expression. As an alternative tissue specific promoter/enhancers can be used to limit expression to the cell types.

Example 5 – Ribozyme

5

This Example is carried out following the methodology of Example 1. A ribozyme which targets a gene on the biosynthetic pathway that generates melanin is delivered using ELAV. This approach facilitates the identification of transgenics.

10 Example 6 - Use of ELAV for transgenic models of Parkinson's.

Parkinson's disease (PD) is one of the most common neurodegenerative diseases, affecting almost 2% of the population over 65. The disease is characterised by a movement disorder - parkinsonism - symptoms of which are rigidity, resting tremor and bradykinesia (slowness to initiate and carry out movement) . This results from the loss of neurons in the substantia nigra that produce the neurotransmitter dopamine. The causes of PD are largely unknown, although there are a few rare families in which the disease is inherited. In families with autosomal dominant PD two different missense mutations have been mapped in α -synuclein (Polymeropoulos et al 1997; Kruger et al 1998), which is a small phosphoprotein thought to be involved in synaptic vesicle transport. In the case of autosomal recessive juvenile parkinsonism (AR-JP), which develops in adolescence, Kitada et al (1998) showed the gene responsible to be Parkin, an E3 ubiquitin ligase recently proposed to catalyse the ubiquitination of α -synuclein (Shimura et al 2001). It has therefore been suggested that an inability to degrade α -synuclein results in AR-JP and possibly sporadic PD (Haass and Kahle 2001).

The ELAV vector system is used to deliver one or more of the following to mouse spermatogonial stem cells (Nagano et al 2001):

30

1. ribozyme to Parkin

2. mutant α -synuclein allele

3. ribozyme to tyrosine hydroxylase (enzyme required for dopamine synthesis)

Example 7 - Angiogenesis

5

The hypoxia inducible factor (HIF) is a transcriptional complex that plays a central role in oxygen homeostasis. The alpha subunit of HIF is targeted for degradation under normoxic conditions by the von Hippel-Lindau ubiquitylation complex that recognizes a hydroxylated proline residue in HIF. Steady state levels of the protein are consequently low and the transcriptional complex cannot form. A family of prolyl-4-hydroxylases have recently been described (Epstein et al 2001) whose enzyme activity is modulated by hypoxia, iron chelation and cobaltous ions, fulfilling the requirements for being oxygen sensors that regulate HIF. Suppression of prolyl-4-hydroxylase in cultured *Drosophila melanogaster* cells by RNA interference resulted in elevated expression of a hypoxia-inducible gene under normoxic conditions (Bruick and McKnight 2001).

The ELAV vector system is used to deliver:

- 20 1. A ribozyme to prolyl-4-hydroxylase (or VHL). This may lead to constitutive up-regulation of HIF-1alpha subunits, activation of the HIF complex and overexpression of HIF target genes.
- 25 2. Constitutively active HIF-1 (upregulation of HIF in normoxia) or PHD3 (downregulation of HIF in hypoxia).

to mouse oocytes by injection into the perivitelline space (Chan et al 1998; 2001).

The production and applications of transgenic mouse models in health-related research are well documented. The proposed research will enable the development of models for a broad range of human diseases the generation of which are currently unmet by existing 'knockout' methodology.

Advantages over existing technology include the following:

- 1) Increased efficiency of transgene delivery by lentiviral transduction as compared with non-homologous recombination of injected DNA. Pronuclear injection leads to insertion of large tandem arrays of DNA which are unstable and subject to rearrangements and deletions. Lentiviral transduction generally leads to the stable integration of a limited number of vector copies distributed as discrete cassettes in the chromosomal DNA .
- 2) Reduction in turnaround time compared to current 'knock-out'. To produce mice with homozygous gene deletions is a relatively labour-intensive and time-consuming process requiring the cross-breeding of mosaic heterozygotes in which the engineered gene deletion has 'gone germline'. In contrast, by transducing oocytes prior to fertilisation, every cell will contain the ablation cassette. The need for cross-breeding is by-passed resulting in shorter turnaround times and a substantial decrease in the overall number of animals required.
- 3) Flexibility of gene product knock-down. As discussed this technology will be of particular value in establishing disease models where deletion of the gene of interest is lethal. It will be advantageous in all studies where ablation of gene expression is desired at particular developmental stages or restricted to specific tissues.
- 4) HIV vectors have a number of significant disadvantages which may limit their therapeutic application to certain diseases. HIV-1 has the disadvantage of being a human pathogen carrying potentially oncogenic proteins and sequences. There is the risk that introduction of vector particles produced in packaging cells which express HIV gag-pol will introduce these proteins into an individual leading to seroconversion. The present non-primate lentiviral-based vectors used in embodiment of the present invention do not introduce HIV proteins into individuals

Example 8 Production of transgenic avians as bioreactors for the production of proteins.



An EIAV vector encoding for bacterial β -galactosidase, initially 10 microlitres of pONY8Z 5'cppt, is injected directly below the blastoderm stage embryo in new-laid eggs using published technology such as is described in US 5,258,307 or earlier stage embryos using techniques such as those described in WO 90/13626. An inert dye is
5 used for blastoderm stage injections to ensure accurate delivery of the vector. Transduction efficiency is analysed by harvesting embryos at different stages of development during incubation and after hatching. The embryos are then sectioned and stained for β -galactosidase activity to identify which organs are transduced and whether germ cells in the embryonic gonad were transduced. Samples of tissues such
10 as blood and CAM are taken from the embryos and the percentage of cells transduced will be assessed using quantitative PCR. Some of the male embryos are grown to sexual maturity and semen samples will be taken. The likelihood of such birds passing the transgene on to any offspring is assessed using quantitative PCR. The semen is then used to inseminate hens and the embryos are harvested and assessed by
15 quantitative PCR to determine the percentage of transgenic offspring. In addition the level of β -galactosidase expression from the vector in the G₁ population is assessed by staining using X-Gal.

Example 9 Vectors for use in RNA interference (RNAi)

20

Figure 20 illustrates a number of expression cassettes for RNAi which can be used in lentiviruses, for example EIAV) to express siRNA in transgenic cells and animals. In each of these examples, an RNA polymerase III promoter (U6) has been utilised. RNA polymerase III makes a variety of very small, stable RNAs including the small 5S
25 ribosomal RNA and the transfer RNAs. Effective RNAi is mediated by either expression of a short hairpin from a single U6 promoter (Fig.20A) or incorporating two U6 promoters, one expressing a region of sense and the other the reverse complement of the same sequence of the target (Fig. 20B). In the embodiment shown in Figure 20C, two opposing promoters are used to transcribe the sense and antisense
30 regions of the target from the forward and complementary strands of the expression cassette.

Example 10 - Aptazymes for regulating production of short RNAs

The use of viral vectors, e.g. lentiviral vectors for generating transgenics to deliver siRNAs which target a gene product with an important or essential function may result in death of the transgenic animal during development. The ability to regulate transcription of the siRNAs, thereby allowing the silencing effect to be modulated, would be greatly desirable. In this example, we describe the use of an aptamer/ribozyme hybrid ('aptazyme') for regulating the production of functional siRNAs.

Aptamers are nucleic acid molecules which form structures which are able to bind a number of ligands including proteins and drug molecules. By replacing one helix of a hammerhead ribozyme with an aptamer it has been possible to create a catalytic RNA which is able to cleave a substrate (which may be itself) as the result of conformational change induced by the presence or absence of a ligand. Figure 21A illustrates the design of an expression cassette which may be used in vectors of the invention and methods of the invention. In this cassette, an aptazyme is added 5' of siRNA encoded by a short hairpin. This allows the regulated induction (or inhibition) of self-cleavage of the transcript separating the hairpin from the aptazyme structure and hence activating silencing. As illustrated in Figure 21B, addition/removal of a ligand triggers catalytic activity, cleaving the transcript and allowing release of the short hairpin to induce silencing of a target sequence.

Example 11 - Hypoxically induced silencing of VEGF by siRNAs

In this example, we describe the use of an 'aptazyme' for regulating the production of functional siRNAs, in which the protein ligand specific for the aptamer is expressed from the same vector. A vector is constructed as illustrated in Figure 22. The RNA polymerase III U6snRNA gene promoter is used to drive expression of an aptazyme-linked short hairpin against VEGF. Under hypoxic conditions expression from the hypoxic response element (HRE) is induced transcribing gene X which codes for a protein X which is a ligand for the aptamer. Binding of the ligand to the aptazyme

triggers catalysis, release of the short hairpin and consequently gene silencing of vascular endothelial growth factor (VEGF)

Thus VEGF is specifically down-regulated in hypoxia which may be therapeutically
5 beneficial in a number of diseases including proliferative diabetic retinopathy.

In an alternative embodiment, the ligand for the aptamer may be VEGF itself.

**Example 12 - Use of RNA polymerase II promoters for transcribing siRNA
10 precursors.**

In this example, we describe the transcription of siRNA precursors under the control of RNA polymerase II promoters. This achieved by flanking the short hairpin (Figure 23A), or siRNA sequence (Figure 23B) with sequence which codes for, or is a target
15 of, a catalytic RNA such as an aptazyme. Cleavage of the flanking sequences releases the siRNA or short hairpin from the precursor.

In the expression cassette shown in Figure 23A, expression of the short hairpin is under the control of an RNA polymerase II promoter, CMV. Two copies of Tet
20 operator downstream provide an additional level of regulation. Transcription is inhibited in the presence of the Tet repressor protein (in the absence of doxycycline) which may be expressed separately or from the same vector. The transcript is flanked by aptazymes which can be activated to cleave at sites designed to release the short hairpin such that it can initiate gene silencing of the target. In this particular
25 embodiment, it is necessary to use aptazymes rather than ribozymes as the latter would result in autologous cleavage of the lentiviral genome.

In the expression cassette shown in Figure 23B, only expression of the antisense siRNA is under the control of aptazyme regulation. Again the target sequence is
30 flanked by aptazymes which cleave at sites releasing the appropriate RNA sequence to form a duplex with the sense RNA which is constitutively expressed from the U6

promoter. Gene silencing of the target can therefore be switched on or off depending on the presence/absence of the aptazyme ligand.

Example 13 - Use of vectors comprising aptazyme sequences for regulating expression of transgenes

Aptazymes may be used for post-transcriptional regulation of any nucleic acid sequence including genes. The aptazyme is activated (or inhibited) by the addition/removal of the appropriate ligand inducing cleavage of the transcript, either removing part of the transcript, for example the codon for the initiator methionine, or a UTR preventing capping and/or polyadenylation of the transcript. This provides a means of shutting off synthesis of a gene product, for example a therapeutic gene such as Factor IX, if levels are too high. Expression of the transgene can be engineered to be self-regulating in this way.

Figure 24A illustrates a construct for use in modulation of expression of insulin. The aptazyme whose activity is modulated by glucose binding is designed such that high level expression of insulin occurs only when blood glucose levels are high. If glucose fall below a threshold level than the aptazyme is active and the insulin transgene transcript destroyed.

Figure 24B illustrates a construct for use in modulation of expression of Factor IX. In this construct, an aptazyme which is regulated by doxycycline is used to regulate the expression of transgenes / short RNAs both in vitro and in vivo by the administration of doxycycline. In this construct, Tet operator sequences are inserted downstream of the promoter. Transcription is repressed in the presence of the Tet repressor protein (which may be expressed from the same vector) when doxycycline is removed, thereby preventing de novo transcription. As the aptazyme is active in the absence of doxycycline any existing transcripts will be cleaved and degraded.

The T-RexTM (InVitrogen) system could optionally be incorporated in such a strategy to add an additional level of control.

Example 14 - Use of vectors comprising aptazyme sequences for regulating expression of transgenes_– Measures to prevent self-cleavage of vector genome RNA

5

Although production of viral vectors of the invention should be carried out under conditions which should minimise activity of the aptazyme, and hence unwanted destruction of the vector genome by self-cleavage, a preferred measure is to physically separate the aptazyme (or ribozyme) by configuring the vector as a split intron vector (Ismail et al 2000). This ensures that the full sequence of the ribozyme is only present in the transcript encoded by the provirus and not in the RNA genome present in the vector particle.

Figure 25A illustrates the split intron strategy with Figure 28B illustrating suitable vector of this aspect of the invention. During the process of reverse transcription a splice donor, and some of the 5' sequence of the ribozyme, is copied to the 5' viral LTR such that the ribozyme is created only following splicing of the transcribed provirus. Using the specific example shown in Figure 25A, the sequence coding for the aptazyme would be split apart in the genome packaged by viral producer cells such that the region indicated in blue (AGAUCAU) would not be present upstream of the sequence shown in black (GAUGCU). Instead it will be present in the 3' LTR along with additional sequence comprising a splice donor (underlined). Upon reverse transcription this will be copied to the 5' LTR such that it is now upstream of a splice acceptor adjacent to the rest of the aptazyme sequence. Upon transcription the intron sequence, underlined:

25

(GUAAAUAAAACTGGGCTTGTCGAGACAGAGAAGACTCTTGCGTTTCTGAT AGGCACCTATTGGTCTTACTGACATCCACTTTGCCTTTCTCTCCACAG), will be spliced thereby forming a complete aptazyme. Therefore the aptazyme is only present in transduced cells preventing any prior cleavage of the vector genome which would lead to loss of titre.

30

Figure 25B illustrates a split intron vector which may be used in this aspect of the invention. The vector has an EIAV / MLV hybrid LTR. This also has a splice donor inserted downstream of the initiation of transcription and upstream of the EIAV repeat and which contains sequence of the 5' portion of the aptazyme. During reverse
5 transcription the modified 3' LTR is copied to the 5' LTR. Following transcription and splicing the functional aptazyme is created. Activation of the aptazyme cleaves the transcript resulting in its degradation.

An additional means of preventing formation of a potentially active aptazyme within
10 the viral RNA genome is to include sequence at the 3' end of the promoter which is able to base-pair with a part of the aptazyme reducing the possibility of it adopting the correct configuration. This is illustrated in Figure 26. As shown, the 3' end of the U6 promoter has been modified to incorporate sequence which will base pair with the the 5' region of helix I forming a hairpin which will prevent the aptazyme from adopting
15 the configuration necessary for catalytic activity. This will only occur in the RNA genome and not in the transcript as initiation of transcription will be downstream of the sequence modified in the promoter. This would not interfere with the tertiary structure of the transcribed provirus as promoter sequences would not be present.

20 Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as
25 claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in chemistry or biology or related fields are intended to be covered by the present invention. All publications mentioned in the above specification are herein incorporated by reference.

References

- Nagano M, Brinster CJ, Orwig KE, Ryu BY, Avarbock MR, Brinster RL.
Proc Natl Acad Sci U S A. 2001 Nov 6;98(23):13090-5.
- 5 Chan AW, Homan EJ, Ballou LU, Burns JC, Bremel RD.
Proc Natl Acad Sci U S A. 1998 Nov 24;95(24):14028-33.
- Chan AW, Chong KY, Martinovich C, Simerly C, Schatten G.
Science. 2001 Jan 12;291(5502):226
- 10 Vilotte JL, L'Huillier P, Mercier JC.
J Mammary Gland Biol Neoplasia 1998 Jul;3(3):351-362
- Polymeropoulos et al (1997) Science 276(5321):2045-7
- 15 Kruger et al (1998) Nat Genet 18(2):106-8
- Kitada et al (1998) Nature 392(6676):605-8
- 20 Shimura et al (2001) Science 293(5528):239-9
- Haas and Kahle (2001) Science 293(5528):224-5
- Epstein et al (2001) Cell 107(1):43-54
- 25 Bruick and McKnight (2001) Science 294(5545):1337-40
- Soukup and Breaker (1999) Structure Fold Des. Jul 15;7(7):783-91
- 30 Van Deutekom et al (2001) Hum Mol Genet Jul 15;10(15):1547-54
- Vickers et al (2001) Nucleic Acids Res. Mar 15;29(6):1293-9

Lagos-Quintana et al (2001) Science Oct 26; 294 (5543):853-8

Ohkawa et al (2000) Hum Gene Ther. Mar 1;11(4):577-85

- 5 Stetor SR, Rausch JW, Guo MJ, Burnham JP, Boone LR, Waring MJ, Le Grice SF.
Characterization of (+) strand initiation and termination sequences located at the center
of the equine infectious anemia virus genome.
Biochemistry 1999 Mar 23;38(12):3656-67
- 10 Donello JE, Loeb JE, Hope TJ.
Woodchuck hepatitis virus contains a tripartite posttranscriptional regulatory element.
J Virol 1998 Jun;72(6):5085-92
- 15 Mitrophanous K, Yoon S, Rohll J, Patil D, Wilkes F, Kim V, Kingsman S, Kingsman
A, Mazarakis N.
Stable gene transfer to the nervous system using a non-primate lentiviral vector.
Gene Ther 1999 Nov;6(11):1808-18

CLAIMS

1. A method of producing a transgenic cell comprising introducing into a cell a
5 non-primate lentiviral expression vector comprising a nucleotide of interest (NOI).
2. A method according to claim 1 wherein the NOI encodes and is capable of
expressing a therapeutic protein or an aptazyme, or is capable of generating an
antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA
10 or a group 1 intron.
3. A method according to claim 1 or 2 wherein the non-primate lentiviral
expression vector is derived from EIAV, FIV, BIV, CAEV or MVV.
- 15 4. A method of producing a transgenic cell comprising introducing into a cell a
lentiviral expression vector comprising a NOI or an aptazyme, capable of generating
an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-
RNA or a group 1 intron.
- 20 5. A method according to claim 4 wherein the lentiviral expression vector is
derived from EIAV, FIV, BIV, CAEV, MVV or HIV.
6. A method according to any preceding claim wherein the expression vector is
introduced in vivo or ex vivo.
25
7. A method according to claim 6 wherein the cell is in utero.
8. A method according to claim 7 wherein the cell is a perinatal cell.
- 30 9. A method according to claim 8 wherein the cell is an embryonic cell.
10. A method according to claim 9 wherein the cell is a fetal cell.

11. A method according to any preceding claim wherein the cell is capable of giving rise to a germ line change.
- 5 12. A method according to claim 11 wherein the cell is a germ cell.
13. A method according to claim 11 wherein the cell is involved in gametogenesis.
14. A method according to any one of claims 11 to 13 wherein the cell is an
10 oocyte, an oviduct cell, an ovarian cell, an ovum, an oogonium, a zygote, an ES cell, a blastocyte, a spermatocyte, a spermatid, a spermatozoa, or a spermatogonia.
15. A method according to any preceding claim wherein the lentiviral expression vector is introduced into the cell via the blastoderm, umbilical cord, placenta, or
15 amniotic fluid, uterus, gonads, or by intraperitoneal, intramuscular, intraspinal, intracranial, intravenous, intra-respiratory, gastrointestinal, or intrahepatic administration.
16. A method according to claim 15 wherein the lentiviral expression vector is
20 introduced into a cell in utero via the blastoderm, umbilical cord, placenta, or amniotic fluid, or by intraperitoneal, intramuscular, intraspinal, intracranial, intravenous, intra-respiratory, gastrointestinal, or intrahepatic administration.
17. A method of producing a transgenic cell comprising introducing into a non-
25 dividing cell a lentiviral expression vector comprising an NOI.
18. A method according to claim 17 wherein the lentiviral expression vector is derived from ELAV, FIV, BIV, CAEV, MVV or HIV.
- 30 19. A method according to claim 17 or 18 wherein the NOI encodes and is capable of expressing a protein or an aptazyme, or is capable of generating an antisense

oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA or a group
1 intron.

20. A method according to any one of claims 17 to 19 wherein the cell is capable
5 of giving rise to a germ line change.

21. A method according to claim 20 wherein the cell is a germ cell.

22. A method according to claim 20 wherein the cell is involved in gametogenesis.

10

23. A method according to claim 22 wherein the cell is an oocyte.

24. A method according to any preceding claim wherein the cell is from an animal,
or a yeast.

15

25. A method according to claim 24 wherein the cell is from a non-human
organism.

26. A method according to claim 24 wherein the cell is mammalian.

20

27. A method according to claim 24 wherein the cell is a murine, human, porcine,
bovine, simian, ovine, equine, avian, insect or reptile or piscine cell.

28. A method according to claim 24 wherein the cell is from *C. elegans* or
25 *drosophila*.

29. A method according to any preceding claim wherein the lentiviral expression
vector is pseudotyped.

30. A method according to any preceding claim wherein the lentiviral expression
vector does not contain any functional accessory genes.

31. A method according to any preceding claim wherein the NOI is operably linked to a constitutive, tissue-specific or an inducible promoter.
32. A transgenic cell produced by the method of any preceding claim.
- 5 33. A transgenic organism which is generated from or obtainable by generation from a transgenic cell according to claim 32 or from the method as defined in any one of claims 1 to 31.
- 10 34. A transgenic organism according to claim 33 wherein the NOI is expressed in an oviduct cell, reproductive tract cell, haematopoietic cell, (including monocytes, macrophages, lymphocytes, granulocytes, or progenitor cells of any of these); secretory cell, mammary cell, endothelial cell, tumour cell, stromal cell, astrocyte, or glial cell, muscle cell, epithelial cell, neuron, fibroblast, hepatocyte, kidney, liver,
- 15 heart or lung cell.
35. A transgenic organism according to claim 33 or 34 wherein the organism is avian.
- 20 36. A transgenic organism according to claim 35 wherein the organism is a fowl such as a chicken, duck or goose.
37. An transgenic egg derived from a transgenic organism according to any one of claims 33 to 36.
- 25 38. A transgenic organism or egg according to any one of claims comprising at least one NOI which encodes and is capable of expressing a protein.
39. A transgenic organism or egg according to claim 38 further comprising at least
- 30 one NOI which is capable of generating an aptazyme, an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA or a group 1 intron.

40. A vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises:

- (a) a second nucleotide sequence encoding an aptazyme; and
- 5 (b) a third nucleotide sequence capable of generating a polynucleotide;

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.

10

41. A vector according to claim 40 wherein said polynucleotide is an RNA molecule capable of modulating expression of a target gene.

42. A vector according to claim 41 wherein the RNA molecule is selected from the group comprising an aptazyme, siRNA, short hairpin RNA, microRNA, anti-sense RNA and a ribozyme.

15

43. A vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises:

20

- (a) a second nucleotide sequence encoding an aptazyme; and
- (b) a third nucleotide sequence comprising a NOI;

wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave the transcript of the first nucleotide sequence such that expression of said NOI is inhibited.

25

44. A vector according to claim 43 wherein the aptazyme is activatable to cleave the transcript of the first nucleotide sequence at a position within the transcript of the third nucleotide sequence.

30

45. A vector according to claims 43 or 44 wherein the NOI encodes a therapeutic protein.
46. A vector according to any one of claims 40 to 45 wherein the aptazyme is
5 activated by a ligand.
47. A vector according to any one of claims 40 to 45 wherein the aptazyme is deactivated by a ligand.
- 10 48. A vector according to claim 46 or 47 wherein the vector comprises a fourth nucleotide sequence encoding the ligand of claims 46 or 47.
49. A vector according to 48 wherein the nucleotide sequence encoding the ligand is operatively linked to a promoter.
- 15 50. A vector according to claims 46 to 49 wherein the ligand is selected from the group comprising polypeptides and fragments thereof, linear peptides, cyclic peptides, and nucleic acids which encode therefor, synthetic and natural compounds including low molecular weight organic or inorganic compounds and antibodies.
51. A vector according to claims 46 to 49 wherein the ligand is selected from the group comprising FMN, doxycycline and VEGF, tetracycline and glucose.
- 20 52. A vector according to any one of claims 40 to 51 wherein (a) and (b) are operably linked to a promoter.
53. A vector according to claim 52 wherein the promoter is selected from the group comprising RNA polymerase III promoter and RNA polymerase II promoter.
- 25 54. A vector according to claim 52 wherein the promoter is operably linked to at least one copy of a tetracycline responsive element (TRE) such that transcription of the

first nucleotide sequence is regulated by a tetracycline modulator and tetracycline or derivative thereof.

55. A vector according to claim 54 wherein the vector comprises a fifth nucleotide
5 sequence encoding a tetracycline modulator.

56. A vector according to any one of claims 52 to 55 wherein the promoter
contains a sequence at its 3' end which is able to base-pair with a part of the aptazyme
such as to form a hairpin to prevent formation of active aptazyme within the viral RNA
10 genome.

57. A vector according to any one of claims 40 to 56 in the form of a viral vector.

58. A vector according to claim 57 wherein the vector is configured as a split
15 intron vector such as to prevent formation of active aptazyme within the viral RNA
genome.

59. A vector according to claim 57 or 58 wherein the vector system is derived from
a retrovirus, a lentivirus, an adenovirus, an adeno-associated vector, a herpes vector, a
20 pox viral vector, a parvovirus vector and a baculoviral vector.

60. A method of producing a transgenic cell using a vector according to any one of
claims 40 to 59.

25 61. A transgenic organism which is generated from or obtainable by generation
from a transgenic cell as defined in claim 60.

62. A method according to any one of claims 1 to 31 using the viral vector of any
one of claims 46 to 59.

30


63. A transgenic cell produced by the method of claim 62.

64. A transgenic organism which is generated from or obtainable by generation from a transgenic cell of claim 63.

5 65. A transgenic organism according to claim 64 wherein the NOI is expressed in an oviduct cell, reproductive tract cell, albumin, haematopoietic cell, (including monocytes, macrophages, lymphocytes, granulocytes, or progenitor cells of any of these); secretory cell, mammary cell, endothelial cell, tumour cell, stromal cell, or glial cell, muscle cell, epithelial cell, neuron, fibroblast, hepatocyte, astrocyte, kidney, liver, heart or lung cell.

Fig. 1


Liver tissue and histology after in utero injection of ElAV lentivirus



Foetal intravenous injection of ElAV viral vector expression LacZ


FIG 2

Tissue and histology after in utero injection of EIAV lentivirus



Foetal intravenous injection of EIAV viral vector expression LacZ

Foetal Intraplasmal Delivery - Dorsal Root Ganglia




Seven days post foetal intraspinal injection of EIAV viral vector
expression nuclear localising LacZ

Figure 3

Figure 4

Foetal Intravenous Delivery - Dorsal Root Ganglia




**Seven days post foetal intravenous injection of EIAV viral vector
expression nuclear localising LacZ**

Figure 5


Foetal Intravenous Delivery

- Liver



Seven days post foetal intravenous injection of ElAV viral vector
expression nuclear localising LacZ

Foetal Intravenous Delivery - Renal Glomeruli




Seven days post foetal intravenous injection of ELAV viral vector
expression nuclear localising LacZ

Figure 6

Figure 7

Foetal Intravenous Delivery - Pancreas



**Seven days post foetal intravenous injection of ElAV viral vector
expression nuclear localising LacZ**

Figure 8

Foetal Intramuscular Delivery - Skeletal Muscle




**Seven days post foetal intramuscular injection of EIAV viral vector
expression nuclear localising LacZ**

Figure 9

Foetal Intrapерitoneal Delivery - Skeletal Muscle: Diaphragm

Intrapерitoneal injection, diaphragm	Planar histology, diaphragm	Transverse section histology, diaphragm
---	--------------------------------	--



Two weeks post foetal intraperitoneal injection of EIAV viral vector
expression LacZ

Figure 10

Foetal Intramuscular Delivery - Skeletal Muscle

Intramuscular injection,
leg



Planar histology,
leg muscle




Transverse section
histology, diaphragm



Two weeks post foetal intramuscular I injection of EIAV viral vector
expression LacZ

Figure 11



X-Gal visulation for β -galactosidase 96 hours after intra-thoracic and intra-peritoneal EIAV lentivirus. The left-hand figure shows a sagittal section with the viscera removed. The diaphragm has been excised and is viewed anteriorly in the right hand figure.

Fig 12

Schematic of ELAV genomes, with sizes.
These are used for transfection. Upon transfection the
3' LTR will be copied to the 5' LTR


Size (kb)

pONY8.1 G



2.6

pONY8.4ZCG



6.1

pONY8.GCZ



6.1



CMV Promoter



ELAV U3



ELAV R region



ELAV U5 region



Packaging signal



Modified packaging signal



Hybrid U3

AGATCTTGAATAATAAAATGTGTGTTTGTCCGAAATACGCGTTTTGAGATTTCTGTGCGC
GACTAAATTCATGTCGCGCGATAGTGGTGTATCGCCGATAGAGATGGCGATATTGGAA
AAATTGATATTTGAAAATATGGCATATTGAAAATGTCGCCGATGTGAGTTTCTGTGTAAC
TGATATCGCCATTTTCCAAAAGTGATTTTGGGCATACGCGATATCTGGCGATAGCGCT
TATATCGTTTACGGGGGATGGCGATAGACGACTTTGGTGACTTGGGCGATTCTGTGTGTC
GCAAATATCGCAGTTTCGATATAGGTGACAGACGATATGAGGCTATATCGCCGATAGAGG
CGACATCAAGCTGGCACATGGCCAATGCATATCGATCTATACATTGAATCAATATTGGCC
ATTAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCA
TACGTTGTATCCATATCGTAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCC
ATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCA
TAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCTGGCTGACC
GCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAAT
AGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGT
ACATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCC
CGCCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTCCTACTTGGCAGTACATCTA
CGTATTAGTCATCGCTATTACCATGGTGATTGCGGTTTTGGCAGTACACCAATGGGCGTGG
ATAGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTT
GTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACACTGCGATCGCCCGCC
CCGTTGACGCAAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGT
TTAGTGAACCGGGCACTCAGATTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGAAAAGG
CCTTTGTAATAAATATAATTCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTTTCGAGATC
CTACAGTTGGCGCCCGAACAGGGACCTGAGAGGGGGCGCAGACCCTACCTGTTGAACCTGG
CTGATCGTAGGATCCCCGGGACAGCAGAGGAGAACTTACAGAAGTCTTCTGGAGGTGTTT
CTGGCCAGAACACAGGAGGACAGGTAAGATTGGGAGACCCTTTGACATTGGAGCAAGGC
G
CTCAAGAAGTTAGAGAAGGTGACGGTACAAGGGTCTCAGAAATTAACACTACTGGTAACTGT
AATTGGGCGCTAAGTCTAGTAGACTTATTTTCATGATACCAACTTTGTAAAAGAAAAGGAC
TGGCAGCTGAGGGATGTCATTCCATTGCTGGAAGATGTAACCTCAGACGCTGTCAGGACAA
GAAAGAGAGGCCTTTGAAAGAACATGGTGGGCAATTTCTGCTGTAAAGATGGGCCTCCAG
ATTAATAATGTAGTAGATGGAAAGGCATCATTCCAGCTCCTAAGAGCGAAATATGAAAAG
AAGACTGCTAATAAAAAGCAGTCTGAGCCCTCTGAAGAATATCTCTAGAAGTGTGGATC
CCCCGGGCTGCAGGAGTGGGGAGGCACGATGGCCGCTTTGGTCGAGGCGGATCCGGCCAT
TAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTCATGTCCAACATTACCGCCAT
CGTTGTATCCATATCATAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCCAT
GTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATA
GCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCTGGCTGACCGC
CCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAG
GGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTAC
ATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCG
CCTGGCATTATGCCCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCTACG
TATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGCGTGGAT
AGCGGTTTGACTCACGGGGATTTCCAAGTCTCCACCCCATTGACGTCAATGGGAGTTTGT
TTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACACTCCGCCCCATTGACGC
AAATGGGCGGTAGGCATGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGAACC
GTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGGACC
GATCCAGCCTCCGCGGCCCAAGCTTGTGGGATCCACCGGTCGCCACCATGGTGAGCAA
GGGCGAGGAGCTGTTACCGGGGTGGTGGCCATCCTGGTCGAGCTGGACGGCGACGTAAA
CGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGAC
CCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCACCCCTCGTGACCAC
CCTGACCTACGGCGTGCAGTGCTTACGCCGCTACCCCGACCACATGAAGCAGCAGCACTT
CTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGA
CGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCAT
CGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGT
A
CAACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGT
GAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCA

FIG 13 cont.

GCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCAC
CCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTT
CGTGACCGCCCGCCGGGATCACTCTCGGCATGGACGAGCTGTACAAGTAAAGCGGCCGCGA
CTCTAGAGTCGACCTGCAGGAATTCGATATCAAGCTTATCGATACCGTCGAATTGGAAGA
GCTTTAAATCCTGGCACATCTCATGTATCAATGCCTCAGTATGTTTAGAAAAACAAGGGG
GGAAGTGTGGGGTTTTATGAGGGGTTTTATAAATGATTATAAGAGTAAAAAGAAAGTTG
CTGATGCTCTCATAACCTTGTATAACCCAAAGGACTAGCTCATGTTGCTAGGCAACTAAA
CCGCAATAACCGCATTGTGACGCGAGTTCCTTGGTGACGCGTTAACTTCCTGTTTT
TACAGTATATAAGTGTGTTGATTCTGACAATTGGGCACTCAGATTCTGCGGTCTGAGTCC
CTTCTCTGCGGCTGAAAAGGCCTTTGTAATAAATATAATTCTCTACTCAGTCCCTGTC
TCTAGTTTGTCTGTTTCGAGATCCTACAGAGCTCATGCCTTGGCGTAATCATGGTCATAGC
TGTTTCCTGTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCA
TAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCT
CACTGCCCCGCTTTCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAAC
GCGCGGGGAGAGGCGGTTTTCGTATTGGGCGCTCTTCCGCTTCCTCGCTCACTGACTCGC
TGCGCTCGGTCTGCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGT
TATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAG
G
CCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACG
AGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGA
T
ACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTA
CCGGATACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCT
GTAGGTATCTCAGTTCGGTGTAGGTCGTTCCGCTCCAAGCTGGGCTGTGTGCACGAACCCC
CCGTTACGCCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAA
GACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATG
TAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAG
TATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTT
GATCCGGCAAAACAAACCACCGCTGGTAGCGGTGTTTTTTGTTTGCAAGCAGCAGATTA
CGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGTCTGACGCTC
AGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCA
CCTAGATCCTTTTAAATTAATAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAA
CTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTAT
TTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCT
TACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATT
TATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTAT
CCGCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTA
ATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTCGTTTG
GTATGGCTTCATTACGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGT
TGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTCAGAAGTAAGTTGGCCG
CAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTATGCCATCCG
TAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGATGC
GGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAA
CTTTAAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTCTCAAGGATCTTAC
CGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGACCCCACTGATCTTCAGCATCTT
TTACTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGG
GAATAAGGGCGACACGGAAATGTTGAATACCTACTCTTCTTTTCAATATTATTGAA
GCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTAGAAAAATA
AACAAATAGGGTTCCGCGCACATTTCCCGGAAAAGTGCCACCTAAATTGTAAGCGTTAA
TATTTTGTAAATTCGCGTTAAATTTTTGTAAATCAGCTCATTTTTTAACCAATAGGC
CGAAATCGGCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTTGAGTGTTGT
TCCAGTTTGAACAAGAGTCCACTATTAAGAAGCTGGACTCCAACGTCAAAGGGCGGAAA
AACCGTCTATCAGGGCGATGGCCCACTACGTGAACCATCACCTAATCAAGTTTTTTGGG
GTCGAGGTGCCGTAAAGCACTAAATCGGAACCCTAAAGGGAGCCCCCGATTTAGAGCTTG
ACGGGGAAAGCCAACCTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCGGC

Fig 14

AGATCTTGAATAATAAAAATGTGTGTTTGTCCGAAATACGCGTTTTGAGATTTCTGTGCGC
GACTAAATTCATGTGCGCGGATAGTGGTGTATTCGCGGATAGAGATGGCGATATTGGAA
AAATTGATATTTGAAAATATGGCATATTGAAAATGTGCGCGATGTGAGTTTTCTGTGTAAC
TGATATCGCCATTTTTCCAAAAGTGATTTTTGGGCATACGCGATATCTGGCGATAGCGCT
TATATCGTTTACGGGGGATGGCGATAGACGACTTTGGTGACTTGGGCGATTCTGTGTGTC
GCAATATCGCAGTTTCGATATAGGTGACAGACGATATGAGGCTATATCGCCGATAGAGG
CGACATCAAGCTGGCACATGGCCAATGCATATCGATCTATACATTGAATCAATATTGGCC
ATTAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCA
TACGTTGTATCCATATCGTAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCC
ATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCA
TAGCCCATATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACC
GCCCAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAAT
AGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGT
ACATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCC
CGCCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTCTACTTGGCAGTACATCTA
CGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTGGCAGTACACCAATGGGCGTGG
ATAGCGGTTTGACTCACGGGGATTTCCAAAGTCTCCACCCCATGACGTCAATGGGAGTTT
GTTTTGGCACCAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTGCGATCGCCCGCC
CCGTTGACGCAAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGCTCGT
TTAGTGAACCGGGCACTCAGATTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGAAAAGG
CCTTTGTAATAAATAAATTCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTTTCGAGATC
CTACAGTTGGCGCCCGAACAGGGACCTGAGAGGGGCGCAGACCCTACCTGTTGAACCTGG
CTGATCGTAGGATCCCCGGGACAGCAGAGGAGAACTTACAGAAGTCTTCTGGAGGTGTT
CTGGCCAGAACACAGGAGGACAGGTAAGATTGGGAGACCCTTTGACATTGGAGCAAGGC
G
CTCAAGAAGTTAGAGAAGGTGACGGTACAAGGGTCTCAGAAATTAATACTACTGGTAACTGT
AATTGGGCGCTAAGTCTAGTAGACTTATTTCAATTGATACCAACTTTGTAAAAGAAAAGGA
CTGGCAGCTGAGGGATTGTCAATCCATTGCTGGAAGATTGTAATCAGACGCTGTCAGGA
CAAGAAAGAGAGGGCCTTTGAAAGAACATTGGTGGGCAATTTCTGCTGTAAAGATTGGGCC
TCCAGATTAATAATTGTAGTAGATTGGAAAGGCATCATTCCAGCTCCTAAGAGCGAAATA
TTGAAAAGAAGACTGCTAATAAAAAAGCAGTCTGAGCCCTCTGAAGAATATCTCTAGAACT
AGTGGATCCCCCGGGCTGCAGGAATTCGATATCAAGCTTCAGCTGCTCGAGGATCTGCGG
ATCCGGGGAATTTCCCACTCTCAGGATCCACCATGGGGGATCCCGTCTGTTTACAAACGTC
GTGACTGGGAAAAACCTGGCGTTACCCAACTTAATCGCCTTGACGACATCCCCCTTTTCG
CCAGCTGGCGTAATAGCGAAGAGGCCCGCACCGATCGCCCTTCCCAACAGTTGCGCAGCC
TGAATGGCGAATGGCGCTTTGCCTGGTTTCCGGCACCAAGCGGTGCCGGAAGCTGGC
TGGAGTGCGATCTTCTGAGGCCGATACTGTGCTGCTCCCTCAAACCTGGCAGATGCACG
GTTACGATGCGCCCATCTACACCAACGTAACCTATCCCATACGGTCAATCCGCCGTTTG
TTCCACGGAGAAATCCGACGGGTTGTTACTCGCTCACATTTAATGTTGATGAAAGCTGGC
TACAGGAAGGCCAGACGCGAATTTTGTATGGCGTTAACTCGGCGTTTCATCTGTGGT
GCAACGGGCGCTGGGTGCGTTACGGCCAGGACAGTCTGTTGCGCTCTGAATTTGACCTGA
GCGCATTTTACGCGCCGGAGAAAACCGCCTCGCGGTGATGGTGCTGCGTTGGAGTGACG
GCAGTTATCTGGAAGATCAGGATATGTGGCGGATGAGCGGCATTTTCCGTGACGTCTCGT
TGCTGCATAAACCGACTACACAAATCAGCGATTTCCATGTTGCCACTCGCTTTAATGATG
ATTTACGCCGCGCTGTACTGGAGGCTGAAGTTCAGATGTGCGGCGAGTTGCGTGACTACC
TACGGGTAAACAGTTTCTTTATGGCAGGGTGAAACCGCAGGTGCCAGCGGCACCGCGCCTT
TCGGCGGTGAAATTATCGATGAGCGTGGTGGTTATGCCGATCGCGTCACACTACGTCTGA
ACGTCGAAAACCCGAAACTGTGGAGCGCCGAAATCCCGAATCTCTATCGTGGCGGTGGTTG
AACTGCACACCGCCGACGGCACGCTGATTGAAGCAGAAGCCTGCGATGTCGGTTTCCGCG
AGGTGCGGATTGAAAATGGTCTGCTGCTGCTGAACGGCAAGCCGTTGCTGATTGAGGCG
TTAACCGTACAGAGCATCATCCTCTGCATGGTCAGGTCATGGATGAGCAGACGATGGTGC
AGGATATCCTGCTGATGAAGCAGAACTTTAACGCCGTGCGCTGTTTCGATTATCCGA
ACCATCCGCTGTGGTACACGCTGTGCGACCGCTACGGCCTGTATGTGGTGGATGAAGCCA
ATATTGAAACCCACGGCATGGTGCCAATGAATCGTCTGACCGATGATCCGCGCTGGCTAC
CGGCGATGAGCGAACGCGTAACGCGAATGGTGCAGCGCGATCGTAATACCCGAGTGTGA
TCATCTGGTCGCTGGGGAATGAATCAGGCCACGGCGCTAATCACGACGCGCTGTATCGCT

FIG 14 C

GGATCAAATCTGTGATCCTTCCCGCCCGGTGCAGTATGAAGGCGGCGGAGCCGACACCA
CGGCCACCGATATTATTTGCCCGATGTACGCGCGCGTGGATGAAGACCAGCCCTTCCCGG
CTGTGCCGAAATGGTCCATCAAAAAATGGCTTTCGCTACCTGGAGAGACGCGCCCGCTGA
TCCTTTGCGAATACGCCACGCGATGGGTAACAGTCTTGGCGGTTTCGCTAAATACTGGC
AGGCGTTTCGTAGTATCCCCGTTTACAGGGCGGCTTCGTCTGGGACTGGGTGGATCAGT
CGCTGATTAAATATGATGAAAACGGCAACCCGTGGTCGGCTTACGGCGGTGATTTTGGCG
ATACGCCGAACGATCGCCAGTTCTGTATGAACGGTCTGGTCTTTGCCGACCGCACGCCGC
ATCCAGCGCTGACGGAAGCAAAACACCAGCAGCAGTTTTTCCAGTTCCGTTTATCCGGGC
AAACCATCGAAGTGACCAGCGAATACCTGTTCGTATAGCGATAACGAGCTCCTGCACT
GGATGGTGGCGCTGGATGGTAAGCCGCTGGCAAGCGGTGAAGTGCCTCTGGATGTGCGCTC
CACAAGGTAAACAGTTGATTGAACTGCCTGAACTACCGCAGCCGAGAGCGCCGGCAAC
TCTGGCTCACAGTACGCGTAGTGCAACCGAACGCGAECGCTGGTCAGAAGCCGGGCACA
TCAGCGCCTGGCAGCAGTGGCGTCTGGCGGAAAACCTCAGTGTGACGCTCCCCGCCGCGT
CCCACGCCATCCCGCATCTGACCACCAGCGAAATGGATTTTTCATCGAGCTGGGTAAATA
AGCGTTGGCAATTTAACCGCCAGTCAGGCTTTCTTTCACAGATGTGGATTGGCGATAAAA
AACAACCTGCTGACGCCGCTGCGCGATCAGTTCACCCGTGCACCGCTGGATAACGACATTG
GCGTAAGTGAAGCGACCCGCATTGACCCTAACGCCTGGGTGCAACGCTGGAAGGCGGCGG
GCCATTACCAGGCCGAAGCAGCGTTGTTGCAAGTGCACGGCAGATACACTTGCTGATGCGG
TGCTGATTACGACCGCTCACGCGTGGCAGCATCAGGGGAAAACCTTATTTATCAGCCGGA
AAACCTACCGGATTGATGGTAGTGGTCAAATGGCGATTACCGTTGATGTTGAAGTGGCGA
GCGATACACCGCATCCGGCGCGGATTGGCCTGAACTGCCAGCTGGCGCAGGTAGCAGAGC
GGGTAAACTGGCTCGGATTAGGGCCGCAAGAAAACCTATCCCGACCGCCTTACTGCCGCT
GTTTTGACCGCTGGGATCTGCCATTGTGACACATGTATACCCCGTACGTCTTCCCGAGCG
AAAACGGTCTGCGCTGCGGGACGCGCGAATTGAATTATGGCCACACCAGTGGCGCGGCG
ACTTCCAGTTCAACATCAGCCGCTACAGTCAACAGCAACTGATGGAAACCAGCCATCGCC
ATCTGCTGCACGCGGAAGAAGGCACATGGCTGAATATCGACGGTTTCCATATGGGGATTG
GTGGCGACGACTCCTGGAGCCCGTCAGTATCGGCGGAATTCAGCTGAGCGCCGGTCGCT
ACCATACCAGTTGGTCTGGTGTCAAAAATAATAAACCAGGCGAGGGGGATCCGCAGA
TCCGGCTGTGGAATGTGTGTGAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCA
GAAGTATGCAAAGCTAGAACTAGTGGATCCCCCGGGCTGCAGGAGTGGGGAGGCACGAT
G
GCCGCTTTGGTTCGAGGCGGATCCGGCCATTAGCCATATTATTCATTGGTTATATAGCATA
AATCAATATTGGCTATTGGCCATTGCATACGTTGTATCCATATCATAATATGTACATTTA
TATTGGCTCATGTCCAACATTACCGCCATGTTGACATTGATTATTGACTAGTTATTAATA
GTAATCAATTACGGGGTCATTAGTTCATAGCCCATATATGGAGTTCGCGGTTACATAACT
TACGGTAAATGGCCCGCTGGCTGACCGCCCAACGACCCCCGCCCATGACGTCAATAAT
GACGTATGTTCCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGAGTA
TTTACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCC
TATTGACGTCAATGACGGTAAATGGCCCGCTGGCATTATGCCAGTACATGACCTTATG
GGACTTTCTACTTGGCAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGTGCG
GTTTTGGCAGTACATCAATGGGCGTGGATAGCGGTTTGAATCACGGGGATTTCCAAAGTCT
CCACCCCATGACGTCAATGGGAGTTTGTGTTTGGCACCAAAATCAACGGGACTTTCCAAA
ATGTCGTAACAACTCCGCCCCATTGACGCAATGGGCGGTAGGCATGTACGGTGGGAGGT
CTATATAAGCAGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTG
TTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCGCGGCCCAAGCTTGTTGG
GATCCACCGGTGCGCCACCATGGTGAGCAAGGGCGAGGAGCTGTTACCGGGGTGGTGGCC
ATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGC
GAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTG
CCCGTCCCTGGCCCAACCTCGTGACCACCTGACCTACGGCGTGCAGTGCTTCAGCCGC
TACCCCGACCATGAAGCAGCAGCACTTCTCAAGTCCGCCATGCCGAAGGCTACGTC
CAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAG
TTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGAC
GGCAACATCCTGGGGCACAAGCTGGAGTACAACATAACAGCCACAACGTCTATATCATG
GCCGACAAGCAGAAGAACGGCATCAAGGTGAACCTCAAGATCCGCCACAACATCGAGGA
C
GGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTG
CTGCTGCCCAGCAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAG
AAGCGCGATCATATGGTCTGCTGGAGTTCGTGACCGCCGCGGGGATCACTCTCGGCATG
GACGAGCTGTACAAGTAAAGCGGCCGCGACTCTAGCCTGCAGGAATTCGATATCAAGCTT
ATCGATACCGTCGAATTGGAAGAGCTTTAAATCCTGGCACATCTCATGTATCAATGCCTC

FIG 14 c

AGTATGTTTAGAAAAACAAGGGGGGAAGTGTGGGGTTTTATGAGGGGTTTTATAAAAAAT
GAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCATTTTGAAGGCATG
GAAAAATACATAACTGAGAATAGAGAAGTTCAGATCAAGGTCAGGAACAGATGGAACAG
C
TGAATATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCTGCCCCGGCTCAGGGCCAAG
AACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCTGCCC
CGGCTCAGGGCCAAGAACAGATGGTCCCCAGATGCGGTCCAGCCCTCAGCAGTTTCTAGA
GAACCATCAGATGTTTCCAGGGTGCCCCAAGGACCTGAAATGACCCTGTGCCTTATTTGA
ACTAACCAATCAGTTCGCTTCTCGCTTCTGTTTCGCGCGCTTCTGCTCCCCGAGCTCAATA
AAAGAGCCCAACCCCTCACTCGGGGGGCACTCAGATTCTGCGGTCTGAGTCCCTTCTC
TGCTGGGCTGAAAAGGCCTTTGTAATAAATAAATTCTCTACTCAGTCCCTGTCTCTAGT
TTGTCTGTTTCGAGATCCTACAGAGCTCATGCCTTGCGTAATCATGGTCATAGCTGTTTC
CTGTGTGAAATTTGTTATCCGCTCACAATTCACACAACATACGAGCCGGAAGCATAAAGT
GTAAAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGC
CCGCTTTCCAGTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGG
GGAGAGGCGGTTTGCGTATTGGGCGCTCTTCCGCTTCTCGCTCACTGACTCGCTGCGCT
CGGTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCA
CAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAG
GA
ACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATC
ACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAAG
G
CGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGAT
ACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCAGCTGTAGGT
ATCTCAGTTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTT
AGCCCGACCGCTGCGCCTTATCCGGTAACATATCGTCTTGAGTCCAACCCGGTAAGACACG
ACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCG
GTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTG
GTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCG
GCAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCA
GAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGA
ACGAAAACCTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGA
TCCTTTTAAATTAATAAATGAAGTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGT
CTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTT
CATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATAAGGGAGGGCTTACCAT
CTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAG
CAATAAACAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTTATCCGCCT
CCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTT
TGCGCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTACGCTCGTCTGTTTGGTATGG
CTTCATTACAGTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCA
AAAAAGCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGTCAGAAGTAAGTTGGCCGAGTGT
TATCACTCATGGTTATGGCAGCACTGCAATAATTCTTACTGTGTCATGCCATCCGTAAGAT
GCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGAC
CGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAA
AAGTGCTCATATTGGAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGCTGT
TGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAAGTATCTTCAGCATCTTTTACTT
TCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATA
A
GGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAGCATTT
ATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTAGAAAAATAAACAA
TAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTAAATTGTAAGCGTTAATATTT
GTTAAAAATTCGCGTTAAATTTTGTAAATCAGCTCATTTTAAACCAATAGGCCGAAAT
CGGCAAAATCCCTTATAAATCAAAAGAATAGACCGAGATAGGGTTGAGTGTGTTCCAGT
TTGGAACAAGAGTCCACTATTAAGAAGCTGGACTCCAACGTCAAAGGGCGAAAAACCGT
CTATCAGGGCGATGGCCCACTACGTGAACCATCACCTAATCAAGTTTTTGGGGTTCGAG
GTGCCGTAAAGCACTAAATCGGAACCCTAAAGGGAGCCCCCGATTTAGAGCTTGACGGG
AAAGCCAACCTGGCTTATCGAAATTAATACGACTCACTATAGGGAGACCGGC

Fig 15

AGATCTTGAATAATAAAATGTGTGTTTGTCCGAAATACGCGTTTTGAGATTTCTGTGCGC
GACTAAATTCATGTCGCGCGATAGTGGTGTATTCGCCGATAGAGATGGCGATATTGGAA
AAATTGATATTTGAAAATATGGCATATTGAAAATGTGCGCGATGTGAGTTTCTGTGTAAC
TGATATCGCCATTTTCCAAAAGTGATTTTGGGCATACGCGATATCTGGCGATAGCGCT
TATATCGTTTACGGGGGATGGCGATAGACGACTTTGGTGACTTGGGCGATTCTGTGTGTC
GCAATATCGCAGTTTCGATATAGGTGACAGACGATATGAGGCTATATCGCCGATAGAGG
CGACATCAAGCTGGCACATGGCCAATGCATATCGATCTATACATTGAATCAATATTGGCC
ATTAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGCA
TACGTTGTATCCATATCGTAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCC
ATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCA
TAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCGCTGGCTGACC
GCCCCAACGACCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAAT
AGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGT
ACATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCC
CGCCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTCTACTTGGCAGTACATCTA
CGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACACCAATGGGCGTGG
ATAGCGGTTTTGACTCACGGGGATTTCGAAGTCTCCACCCCATTGACGTCAATGGGAGTTT
GTTTTGGCACAAAATCAACGGGACTTTCCAAAATGTCGTAACAACTGCGATCGCCCGCC
CCGTTGACGCAAAATGGGCGGTAGGCGGTACGGTGGGAGGTCTATATAAGCAGAGCTCGT
TLAGTGAACCGGGCACTCAGATTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGAAAAGG
CCTTTGTAATAAATATAATTCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTCGAGATC
CTACAGTTGGCGCCCGAACAGGGACCTGAGAGGGGCGCAGACCCTACCTGTTGAACCTGG
CTGATCGTAGGATCCCCGGGACAGCAGAGGAGAACTTACAGAAGTCTTCTGGAGGTGTTT
CTGGCCAGAACACAGGAGGACAGGTAAGATTGGGAGACCCTTTGACATTGGAGCAAGGC
G
CTCAAGAAGTTAGAGAAGGTGACGGTACAAGGGTCTCAGAAATTAACACTACTGGTAACTGT
AATTGGGCGCTAAGTCTAGTAGACTTATTTCAATTGATACCAACTTTGTAAGAAAAGGA
CTGGCAGCTGAGGGATTGTCAATTCATTGCTGGAAGATTGTAACCTCAGACGCTGTCAAG
CAAGAAAGAGAGGGCCTTTGAAAGAACATTGGTGGGCAATTTCTGCTGTAAAGATTGGGC
TCCAGATTAATAATTGTAGTAGATTGGAAAGGCATCATTCCAGCTCCTAAGAGCGAAATA
TTGAAAGAAGACTGCTAATAAAAAGCAGTCTGAGCCCTCTGAAGAATATCTCTAGAGTC
GACGGTACCGCGGGCCCGGGATCCACCGGTGCGCCACCATGGTGAGCAAGGGCGAGGAGC
T
GTTACCGGGGTGGTGCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTT
CAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCAT
CTGACCAACCGGCAAGCTGCCCGTGCCTTGGCCACCTCGTGACCACCTGACCTACGG
CGTGCAGTGCTTCAGCCGCTACCCCGACCATGAAGCAGCAGCACTTCTCAAGTCCGC
CATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTCAAGGACGACGGCAACTACAA
GACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGG
G
CATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACCTACAACA
G
CCACAACGTCTATATCATGGCCGAACAAGCAGAAGAACGGCATCAAGGTGAACTTCAAGAT
CCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCTGAGCACCCAGTCCGCCCT
CATCGGCGACGGCCCCGTGCTGCTGCCCCGACAACCACTACCTGAGCACCCAGTCCGCCCT
GAGCAAAGACCCCAACGAGAAGCGCGATCAGATGGTCCTGCTGGAGTTCTGTGACCGCCGC
CGGATCACTCTCGGCATGGACGAGCTGTACAAGTAAAGCGGCCGCTCTAGAAGTAGTGG
ATCCCCCGGGCTGCAGGAGTGGGAGGACGATGAGGCGCTTTGGTTCGAGGCGGATCCGGC
CATTAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTATTGGCCATTGTC
ATACGTTGTATCCATATCATAATATGTACATTTATATTGGCTCATGTCCAACATTACCGC
CATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTC
ATAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCTGGCTGAC
CGCCCAACGACCCCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAA
TAGGGACTTTCCATTGACGTCAATGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAG

Fig 15a L

TACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAATGGC
CCGCTGGCATTATGCCAGTACATGACCTTATGGGACTTTCCTACTTGGCAGTACATCT
ACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATCAATGGGCGTG
GATAGCGGTTTGACTCACGGGGATTTCGAAGTCTCCACCCATTGACGTCAATGGGAGTT
TGTTTTGGCACCAAAATCAACGGGACTTTCGAATGTCTGTAACAACCTCCGCCCCATTGA
CGAAATGGGCGGTAGGCATGTACGGTGGGAGGTCTATATAAGCAGAGCTCGTTTAGTGA
ACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCATAGAAGACACCGGG
ACCGATCCAGCCTCCGCGGCCCAAGCTTCAGCTGCTCGAGGATCTGCGGATCCGCGGAA
TTCCCCAGTCTCAGGATCCACCATGGGGGATCCCGTCGTTTTACAACGTCGTGACTGGGA
AAACCCTGGCGTTACCCAACCTAATCGCCTTGCAGCACATCCCCCTTTCGCCAGCTGGCG
TAATAGCGAAGAGGGCCCGCACCGATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGA
ATGGCGCTTTGCCTGGTTTCCGGCACCAGAAGCGGTGCCGGAAGCTGGCTGGAGTGCGA
TCTTCTGAGGCCGATACTGTCGTCGTCCCTCAAACCTGGCAGATGCACGGTTACGATGC
GCCATCTACACCAACGTAACCTATCCATTACGGTCAATCCGCGCTTGTTCACCGGA
GAATCCGACGGGTGTTACTCGCTCACATTTAATGTTGATGAAAGCTGGCTACAGGAAGG
CCAGACGCGAATTATTTTTGATGGCGTTAACTCGGCGTTTCATCTGTGGTGCAACGGGCG
CTGGGTGCGTTACGGCCAGGACAGTCGTTTGCCGTCTGAATTTGACCTGAGCGCATTTTT
ACGCGCCGGAGAAAACCGCCTCGCGGTGATGGTGCTGCGTTGGAGTGACGGCATCTATCT
GGAAGATCAGGATATGTGGCGGATGAGCGGCATTTTCCGTGACGTCTCGTGTGTCATAA
ACCGACTACACAAATCAGCGATTTCCATGTTGCCACTCGCTTTAATGATGATTTACGCCG
CGCTGTACTGGAGGCTGAAGTTCAGATGTGCGGCGAGTTGCGTGACTACCTACGGGTAAC
AGTTTCTTTATGGCAGGGTGAAACGCGAGGTGCGCAGCGGCACCGCGCCTTTCGGCGGTGA
AATTATCGATGAGCGTGGTGGTTATGCCGATCGCGTCACACTACGTCTGAACGTCGAAAA
CCCGAACTGTGGAGCGCCGAAATCCCGAATCTCTATCGTGCGGTGGTTGAACTGCACAC
CGCCGACGCGACGCTGATTGAAGCAGAAGCCTGCGATGTCGGTTTCCGCGAGGTGCGGAT
TGAAAATGGTCTGCTGCTGCTGAACGGCAAGCCGTTGCTGATTGAGGCGTTAACCCTCA
CGAGCATCATCCTCTGCATGGTCAGGTCAATGATGAGCAGACGATGGTGAGGATATCCT
GCTGATGAAGCAGAACAACCTTAACGCCGTGCGCTGTTCCGATTATCCGAACCATCCGCT
GTGGTACACGCTGTGCGACCGCTACGGCCTGTATGTGGTGGATGAAGCCAATATTGAAC
CCACGGCATGGTGCCAATGAATCGTCTGACCGATGATCCGCGCTGGCTACCGGCGATGAG
CGAACGCGTAACGCGAATGGTGACGCGGATCGTAATCACCCGAGTGTGATCATCTGGTC
GCTGGGAATGAATCAGGCCACGGCGCTAATCACGACGCGCTGTATCGCTGGATCAAATC
TGTCGATCCTTCCCGCCCGGTGCAGTATGAAGGCGCGGAGCCGACACCACGGCCACCGA
TATTATTTGCCCGATGTACGCGCGCTGGATGAAGACGACCCCTTCCCGGTGTGCCGAA
ATGGTCCATCAAAAAATGGCTTTCGCTACCTGGAGAGACGCGCCCGCTGATCCTTTGCCA
ATACGCCACGCGATGGGTAAACAGTCTTGGCGGTTTCGCTAAATACTGGCAGCGTTTCG
TCAGTATCCCCGTTTACAGGGCGGCTTCGTCTGGGACTGGGTGGATCAGTCGCTGATTAA
ATATGATGAAAACGGCAACCCGTGGTCCGCTTACGGCGGTGATTTTGGCGATACGCCGAA
CGATCGCCAGTTCTGTATGAACGGTCTGGTCTTTGCCGACCGCACGCGCATCCAGCGCT
GACGGAAGCAAAACACCAGCAGCAGTTTTTCCAGTTCCGTTTATCCGGGCAAAACCATCGA
AGTGACCAGCGAATACTGTTCCGTATAGCGATAACGAGCTCCTGCACTGGATGGTGGC
GCTGGATGGTAAGCCGCTGGCAAGCGGTGAAGTGCTCTGGATGTCGCTCCACAAGGTAA
ACAGTTGATTGAACTGCCTGAACTACCGCAGCGGAGAGCGCCGGGCAACTCTGGCTCAC
AGTACGCGTAGTGCAACCGAACGCGACCGCATGGTCAAGAAGCCGGGCACATCAGCGCCTG
GCAGCAGTGGCGTCTGGCGGAAAACCTCAGTGTGACGCTCCCCGCGCGTCCACGCCAT
CCCGCATCTGACCACAGCGAAATGGATTTTGCATCGAGCTGGGTAATAAGCGTTGGCA
ATTTAACCGCCAGTCAGGCTTCTTTACAGATGTGGATTGGCGATAAAAAACAACCTGCT
GACGCCGCTGCGCGATCAGTTCACCCGTGCACCGCTGGATAACGACATTGGCGTAAGTGA
AGCGACCCGATTGACCCTAACGCTGGGTGCAACGCTGGAAGGCGGCGGGCCATTACCA
GGCCGAAGCAGCGTTGTTGAGTGCACGGCAGATACACTTGCTGATGCGGTGCTGATTAC
GACCGCTCACGCGTGGCAGCATCAGGGGAAAACCTTATTTATCAGCCGGAACCTACCG
GATTGATGGTAGTGGTCAAATGGCGATTACCGTTGATGTTGAAGTGGCGAGCGATACACC
GCATCCGGCGCGGATTGGCTGAACTGCCAGTGGCGCAGGTAGCAGAGCGGGTAAACTG
GCTCGGATTAGGGCCGCAAGAAAACCTATCCCGACCGCTTACTGCCGCTGTTTGACCG
CTGGGATCTGCCATTGTCAGACATGTATACCCCGTACGTCTTCCCGAGCGAAAACGCTCT
GCGCTGCGGGACGCGCGAATTGAATTATGGCCACACCAAGTGGCGCGGCGACTTCCAGTT
CAACATCAGCCGCTACAGTCAACAGCAACTGATGGAAACAGCCATCGCCATCTGCTGCA
CGCGGAAGAAGGCACATGGCTGAATATCGACGGTTTCATATGGGGATTGGTGGCGACGA
CTCCTGGAGCCCGTCAGTATCGGCGGAATTCAGCTGAGCGCCGGTCTGCTACCATACCA
GTTGGTCTGGTGTCAAAAATAATAAACCGGGCAGGGGGGATCCGCAGATCCGGCTGTG

Fig. 15

GAATGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTCCCCAGCAGGCAGAAAGTATGCA
AAGCATGCCTGCAGGAATTCGATATCAAGCTTATCGATACCGTCGAATTGGAAGAGCTTT
AAATCCTGGCACATCTCATGTATCAATGCCTCAGTATGTTTAGAAAAACAAGGGGGGAAC
TGTGGGGTTTTTATGAGGGGTTTTATAAAAATGAAAGACCCACCTGTAGGTTTGGCAAG
CTAGCTTAAGTAACGCCATTTTGCAAGGCATGGAAAAATACATAACTGAGAATAGAGAAG
TTCAGATCAAGGTCAGGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTG
GTAAGCAGTTCCTGCCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGCCA
AACAGGATATCTGTGGTAAGCAGTTCCTGCCCCGGCTCAGGGCCAAGAACAGATGGTCCC
CAGATGCGGTCCAGCCCTCAGCAGTTTCTAGAGAACCATCAGATGTTTCCAGGGTGCCCC
AAGGACCTGAAATGACCCTGTGCCTTATTTGAACTAACCAATCAGTTCGCTTCTCGCTTC
TGTTCCGCGCTTCTGCTCCCCGAGCTCAATAAAGAGCCCAACCCCTCACTCGGGGG
GCACTCAGATTCTGCGGTCTGAGTCCCTTCTGCTGGGCTGAAAAGGCCTTTGTAATAA
ATATAATTCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTTTCGAGATCCTACAGAGCTCA
TGCCTTGGCGTAATCATGGTCATAGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAAT
TCCACACAACATACGAGCCGGAAGCATAAAGTGTAAGCCTGGGGTGCCATAGAGTGAG
CTAACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTCCAGTCGGGAAACCTGTCGTG
CCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGGTTTGGGTATTGGGCGCTC
TTCCGCTTCTCGCTCACTGACTCGCTCGCTCGGTCGTTTCGGCTGCGGCGAGCGGTATC
AGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGA
A
CATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGT
T
TTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTG
GCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCG
CTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCGGGAAG
CGTGGCGCTTTCTCATAGCTCAGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTTCGCTC
CAAGCTGGGCTGTGTGCACGAACCCCGCTTCAGCCCCGACCGCTGCGCCTTATCCGGTAA
CTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGACCCACTGG
TAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCCTTGAAGTGGTGCC
TAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTAC
CTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCACCGCTGGTAGCGGTGG
TTTTTTTGTGTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTT
GATCTTTTCTACGGGGTCTGACGCTCAGTGGAAACGAAAACCTCACGTTAAGGGATTTTGGT
CATGAGATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAAAGTATTAA
ATCAATCTAAAGTATATAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGA
GGCACCTATCTCAGCGATCTGTCTATTTTCGTTTACCATAGTTGCCTGACTCCCCGTCGT
GTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCG
AGACCCACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCGA
GCGCAGAAGTGGTCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGA
AGCTAGAGTAAGTAGTTCCGCCAGTTAATAGTTTGCACAACGTTGTTGCCATTGCTACAGG
CATCGTGGTGTACGCTCGTCTGTTTGGTATGGCTTCATTCAGCTCCGGTCCCAACGATC
AAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCCCTCC
GATCGTTGTCAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCA
TAATTCCTTACTGTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAAC
CAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACG
GGATAATACCGCGCCACATAGCAGAACTTTAAAGTGCTCATCATTGGAAAACGTTCTTC
GGGGCGAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCG
TGCACCCAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAAC
AGGAAGGCAAAATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCA
T
ACTCTTCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATA
CATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAA
AGTGCCACCTAAATTGTAAGCGTTAATATTTTGTAAAAATTCGCGTTAAATTTTTGTAA
ATCAGTCAATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAAGAA
TAGACCGAGATAGGGTTGAGTGTTTCCAGTTTGGAAACAAGAGTCCACTATTAAAGAAC
GTGGACTCCAACGTCAAAGGGCGGAAACCGTCTATCAGGGCGATGGCCCACTACGTGAA
CCATCACCCCTAATCAAGTTTTTTGGGGTCGAGGTGCCGTAAAGCACTAAATCGGAACCT
AAAGGGAGCCCCGATTTAGAGCTTGACGGGAAAGCCAACCTGGCTTATCGAAATTAAT
ACGACTCACTATAGGGAGACCGGC

FIGURE 16

Hybrid U3 region, schematic.




FIGURE 17

18

Sequence of hybrid LTR

TGTGGGGTTTTTATGAGGGGTTTTATAATGAAAGACCCACCTGTAGGTTTGGCAAGCT
AGCTTAAGTAACGCCATTTTGCAAGGCATGGAAAAATACATAACTGAGAATAGAGAAGTT
CAGATCAAGGTCAGGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGG
T
AAGCAGTTCCTGCCCCGGCTCAGGGCCAAGAACAGATGGAACAGCTGAATATGGGCCAA
A
CAGGATATCTGTGGTAAGCAGTTCCTGCCCCGGCTCAGGGCCAAGAACAGATGGTCCCCA
GATGCGGTCCAGCCCTCAGCAGTTTCTAGAGAACCATCAGATGTTTCCAGGGTGCCCCAA
GGACCTGAAATGACCCTGTGCCTTATTTGAACTAACCAATCAGTTCGCTTCTCGCTTCTG
TTCGCGCGCTTCTGCTCCCCGAGCTCAATAAAAAGAGCCCACAACCCCTCACTCGGGGGGC
ACTCAGATTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGAAAAGGCCTTTGTAATAAAT

Figure 18 Sequence of pONY8.1ZHyb

AGATCTTGAATAATAAAATGTGTGTTTGTCCGAAATACGCGTTTTGAGATTTCTGTGCGCCG
ACTAAATTCATGTCGCGCGATAGTGGTGTATCGCCGATAGAGATGGCGATATTGGA
AATTGATATTTGAAAATATGGCATATTGAAAATGTCGCCGATGTGAGTTTCTGTGTA
ATATCGCCATTTTTCCAAAAGTGATTTTTGGGCATACGCGATATCTGGCGATAGCGCTTA
TATCGTTTACGGGGGATGGCGATAGACGACTTTGGTGACTTGGGCGATTCTGTGTGTCG
CAAATATCGCAGTTTCGATATAGGTGACAGACGATATGAGGCTATATCGCCGATAGAGG
CGACATCAAGCTGGCACATGGCCAATGCATATCGATCTATACATTGAATCAATATTGGCC
ATTAGCCATATTATTCATTGGTTATATAGCATAAAATCAATATTGGCTATTGGCCATTGCATA
CGTTGTATCCATATCGTAATATGTACATTTATATTGGCTCATGTCCAACATTACCGCCATG
TTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGC
CCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCC
CAACGACCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAG
GGACTTTCATTGACGTCAATGGGTGGAGTATTTACGGTAACTGCCCACTTGGCAGTA
CATCAAGTGTATCATATGCCAAGTCCGCCCCCTATTGACGTCAATGACGGTAAATGGCC
CGCCTGGCATTATGCCCAGTACATGACCTTACGGGACTTTCCTACTTGGCAGTACATCTA
CGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACACCAATGGGCGTG
GATAGCGGTTTGACTCACGGGGATTTCGAAGTCTCCACCCCAATTGACGTCAATGGGAGT
TTGTTTTGGCACCAAATCAACGGGACTTTCAAAATGTCGTAACAACTGCGATCGCCCG
CCCCGTTGACGCAAATGGGCGGTAGGCGTGTACGGTGGGAGGTCTATATAAGCAGAGC
TCGTTTAGTGAACCGGGCACTCAGATTTCTGCGGTCTGAGTCCCTTCTCTGCTGGGCTGA
AAAGGCCTTTGTAAATAAATAAATCTCTACTCAGTCCCTGTCTCTAGTTTGTCTGTTTGA
GATCCTACAGTTGGCGCCCGAACAGGGACCTGAGAGGGGCGCAGACCCTACCTGTTGA
ACCTGGCTGATCGTAGGATCCCCGGGACAGCAGAGGAGAACTTACAGAAGTCTTCTGGA
GGTGTTCTTGCCAGAACACAGGAGGACAGGTAAGATTGGGAGACCCTTTGACATTGGA
GCAAGGCGCTCAAGAAGTTAGAGAAGGTGACGGTACAAGGGTCTCAGAAATTAACACT
GGTAACTGTAATTGGGCGCTAAGTCTAGTAGACTTATTTATGATACCAACTTTGTAAAG
AAAAGGACTGGCAGCTGAGGGATGTCATTCCATTGCTGGAAGATGTAACCTCAGACGCTG
TCAGGACAAGAAAGAGAGGCCCTTTGAAAGAACATGGTGGGCAATTTCTGCTGTAAAGT
GGGCCTCCAGATTAATAATGTAGTAGATGGAAGGCATCATTCCAGCTCCTAAGACGGA
AATATGAAAAGAAAGACTGCTAATAAAAGCAGTCTGAGCCCTCTGAAGAATATCTCTAGA
ACTAGTGGATCCCCCGGGCTGCAGGAGTGGGGAGGCACGATGGCCGCTTTGGTCGAG
GCGGATCCGGCCATTAGCCATATTATTCATTGGTTATATAGCATAAATCAATATTGGCTAT
TGGCCATTGCATACGTTGTATCCATATCATAATATGTACATTTATATTGGCTCATGTCCAA
CATTACCGCCATGTTGACATTGATTATTGACTAGTTATTAATAGTAATCAATTACGGGGTC
ATTAGTTCATAGCCCATATATGGAGTTCGCGGTTACATAACTTACGGTAAATGGCCCGCC
TGGCTGACCGCCCAACGACCCCCCGCCATTGACGTCAATAATGACGTATGTTCCCATAG
TAACGCCAATAGGGACTTTCATTGACGTCAATGGGTGGAGTATTTACGGTAACTGCC
ACTTGGCAGTACATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGACG
GTAATAGCCCCGCTGGCATTATGCCCAGTACATGACCTTATGGGACTTTCCTACTTGG
CAGTACATCTACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTGGCAGTACATC
AATGGGCGTGATAGCGGTTTGACTCACGGGGATTTCGAAGTCTCCACCCCAATTGACGT
CAATGGGAGTTTGTGTTTGGCACCAAATCAACGGGACTTTCAAAATGTCGTAACAACTC
CGCCCCATTGACGCAAATGGGCGGTAGGCATGTACGGTGGGAGGTCTATATAAGCAGA
GCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACCTCCA
TAGAAGACACCGGGACCGATCCAGCCTCCGCGGCCCAAGCTTCAGCTGCTCGAGGAT
CTGCGGATCCGGGGGAATTCCTCCAGTCTCAGGATCCACCATGGGGGATCCCGTCGTTTTA
CAACGTCGTGACTGGGAAAACCTGGCGTTACCCCAACTTAATCGCCTTGACGACATCC
CCCTTTCGCCAGCTGGCGTAATAGCGAAGAGCCCGCACCGATCGCCTTCCCAACAG
TTGCCAGCCCTGAATGGCGAATGGCGCTTTGCTGGTTTCCGGCACCAAGAAGCGGTGC
CGGAAGCTGGCTGGAGTGGGATGCGATCTTCCTGAGGCGGATACTGTCGTGCTCCCTCAAAC
TGGCAGATGCACGGTTACGATGCGCCCATCTACACCAACGTAACCTATCCATTACGGT
CAATCCGCCGTTTGTTCACGGAGAATCCGACGGGTGTTACTCGCTCACATTTAATGT
TGATGAAAGCTGGCTACAGGAAGGCCAGACGCGAATTATTTTTGATGGCGTTAACTCGG
CGTTTCATCTGTGGTGCAACGGGCGCTGGGTGCGTTACGGCCAGGACAGTCGTTTGCC
GTCTGAATTTGACCTGAGCGCATTTTTACGCGCCGGAGAAAACCGCCTCGCGGTGATGG
TGCTGCGTTGGAGTGACGGCAGTTATCTGGAAGATCAGGATATGTGGCGGATGAGCGG
CATTTTCCGTGACGTCTCGTTGCTGCATAAACCAGCTACACAAATCAGCGATTTCCATGT
TGCCACTCGCTTAATGATGATTTACGCCGCGCTGTACTGGAGGCTGAAGTTCAGATGT

Fig 18 cont.

GCGGCGAGTTGCGTGACTACCTACGGGTAACAGTTTCTTTATGGCAGGGTGAAACGCAG
GTCGCCAGCGGCACCGCGCCTTTCCGGCGGTGAAATTATCGATGAGCGTGGTGGTTATG
CCGATCGCGTCACACTACGTCTGAACGTCGAAAACCCGAAACTGTGGAGCGCCGAAATC
CCGAATCTCTATCGTGCGGTGGTTGAACTGCACACCGCCGACGGCAGCGTGATTGAAG
CAGAAGCCTGCGATGTCGGTTTCCGCGAGGTGCGGATTGAAAATGGTCTGCTGCTGCT
GAACGGCAAGCCGTTGCTGATTCGAGGCGTTAACCGTCACGAGCATCATCCTCTGCATG
GTCAGGTCATGGATGAGCAGACGATGGTGCAGGATATCCTGCTGATGAAGCAGAACAAAC
TTTAACGCCGTGCGCTGTTCCGATTATCCGAACCATCCGCTGTGGTACACGCTGTGCGA
CCGCTACGGCCTGTATGTGGTGGATGAAGCCAATATTGAAACCCACGGCATGGTGCCAA
TGAATCGTCTGACCGATGATCCGCGCTGGCTACCGGCGATGAGCGAACGCGTAACGCG
AATGGTGCAGCGCGATCGTAATCACCCGAGTGTGATCATCTGGTCGCTGGGGAATGAAT
CAGGCCACGGCGCTAATCACGACGCGCTGTATCGCTGGATCAAATCTGTGATCCTTCC
CGCCCGGTGCAGTATGAAGGCGGCGGACACCGACCGCCACCGATATTATTTGCC
CGATGTACGCGCGCTGGATGAAGACGAGCCCTTCCCGGCTGTGCCGAAATGGTCCAT
CAAAAAATGGCTTTCGCTACCTGGAGAGACGCGCCCGCTGATCCTTTGCGAATACGCCC
ACGCGATGGGTAACAGTCTTGGCGGTTTCGCTAAATACTGGCAGGCGTTTCGTCAGTAT
CCCCGTTTACAGGGCGGCTTCGTCTGGGACTGGGTGGATCAGTCGCTGATTAAATATGA
TGAAAACGGCAACCCGTTGGTCGGCTTACGGCGGTGATTTTGGCGATACGCCGAACGAT.
CGCCAGTTCTGTATGAACGGTCTGGTCTTTGCCGACCGCACGCCGCATCCAGCGCTGA
CGGAAGCAAAACACCAGCAGCAGTTTTTCCAGTTCCGTTTATCCGGGCAACCATCGAA
GTGACCAGCGAATACCTGTTCCGTCATAGCGATAACGAGCTCCTGCACTGGATGGTGGC
GCTGGATGGTAAGCCGCTGGCAAGCGGTGAAGTGCCCTCTGGATGTCGCTCCACAAGGT
AAACAGTTGATTGAACTGCCTGAACTACCGCAGCCGGAGAGCGCCGGGCAACTCTGGC
TCACAGTACGCGTAGTGCAACCGAACGCGACCGCATGGTCAGAAGCCGGGCACATCAG
CGCCTGGCAGCAGTGGCGTCTGGCGGAAAACCTCAGTGTGACGCTCCCCGCCGCGTCC
CAGCCATCCCGCATCTGACCACCAGCGAAATGGATTTTTGCATCGAGCTGGGTAATAA
GCGTTGGCAATTTAACCGCCAGTCAGGCTTTCTTTCACAGATGTGGATTGGCGATAAAAA
ACAAGTGTGACGCCGCTGCGCGATCAGTTCACCCGTGCACCGCTGGATAACGACATTG
GCGTAAGTGAAGCGACCCGCATTGACCCTAACGCCCTGGGTGCAACGCTGGAAGGCGGC
GGGCCATTACCAGGCCGAAGCAGCGTTGTTGCAGTGCACGGCAGATACACTTGCTGAT
GCGGTGCTGATTACGACCGCTCACGCGTGGCAGCATCAGGGGAAAACCTTATTTATCAG
CCGGAAAACCTACCGGATTGATGGTAGTGGTCAAATGGCGATTACCGTTGATGAAG
TGCGGAGCGGATACACCGCATCCGGCGCGGATTGGCCTGAACTGCCAGCTGGCGCAGGT
AGCAGAGCGGGTAAACTGGCTCGGATTAGGGGCCGCAAGAAAACCTATCCCGACCGCCTT
ACTGCCGCTGTTTTGACCGCTGGGATCTGCCATTGTGACACATGTATACCCCGTACGT
CTTCCCGAGCGAAAACGGTCTGCGCTGCGGGACGCGCGAATTGAATTATGGCCACAC
CAGTGGCGCGGCGACTTCCAGTTCAACATCAGCCGCTACAGTCAACAGCAACTGATGGA
AACCAGCCATCGCCATCTGCTGCACGCGGAAGAAGGCACATGGCTGAATATCGACGGTT
TCCATATGGGGATTGGTGGCGACGACTCCTGGAGCCCGTCAGTATCGGCGGAATTCCA
GCTGAGCGCCGGTCGCTACCATACCAGTTGGTCTGGTGTCAAAAATAATAAACCGG
GCAGGGGGGATCCGCAGATCCGGCTGTGGAATGTGTGTCAGTTAGGGTGTGGAAGTC
CCCAGGCTCCCGCAGGCAGAGTATGCAAAAGCATGCCTGCAGGAATTCGATATCAAG
CTTATCGATACCGTCGAATTGGAAGAGCTTTTAAATCCTGGCACATCTCATGTATCAATGC
CTCAGTATGTTTTAGAAAAACAAGGGGGGAACTGTGGGGTTTTTATGAGGGGTTTTATAAT
GAAAGACCCACCTGTAGGTTTGGCAAGCTAGCTTAAGTAACGCCATTTTGCAAGGCAT
GGAAAAATACATAACTGAGAATAGAGAAGTTCAGATCAAGGTCAGGAACAGATGGAACA
GCTGAATATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCCTGCCCGGGCTCAGGGC
CAAGAACAGATGGAACAGCTGAATATGGGCCAAACAGGATATCTGTGGTAAGCAGTTCC
TGCCCGGGCTCAGGGCCAAGAACAGATGGTCCCCAGATGCGGTCCAGCCCTCAGCAGT
TTCTAGAGAACCATCAGATGTTTCCAGGGTGCCCCAAGGACCTGAAATGACCCTGTGCC
TTATTTGAACTAACCAATCAGTTCGCTTCTCGCTTCTGTTGCGCGCTTCTGCTCCCCGA
GCTCAATAAAAGAGCCCACAACCCCTCACTCGGGGGCACTCAGATTCTGCGGTCTGAG
TCCCTTCTCTGCTGGGCTGAAAAGGCCCTTTGTAAATAATAATTCTCTACTCAGTCCCTG
TCTCTAGTTTGTCTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG
GCTGTTTCTGTGTGAAATTGTTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG
CATAAAGTGTAAAGCCTGGGGTGCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCG
CTCACTGCCCCGCTTTCAGTCGGGAAACCTGTGCTGCCAGCTGCATTAATGAATCGGCC
AACGCGCGGGGAGAGGCGGTTTGCCTATTGGGCGCTTTCGCTTCTCGCTCACTGA
CTCGCTGCGCTCGGTGCTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTA
ATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAGAACATGTGAGCAAAAGGCCA

Fig 18 cont.

GCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGC
CCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAG
GACTATAAAGATACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCG
ACCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTC
TCATAGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTGCTCCAAGCTGGGCT
GTGTGCACGAACCCCCCGTTACGCCCCGACCGCTGCGCCTTATCCGGTAACATCGTCTT
GAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGAT
TAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACG
GCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGA
AAAAGAGTTGGTAGCTCTTGATCCGGCAAAACAAACCACCGCTGGTAGCGGTGGTTTTTT
GTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTT
TCTACGGGGTCTGACGCTCAGTGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAG
ATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAATAAATGAAGTTTTAAATCAATCT
AAAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTA
TCTCAGCGATCTGTCTATTTCTGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAA
CTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCC
ACGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGC
AGAAAGTGGTCCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCT
AGAGTAAGTAGTTGCGCCAGTTAATAGTTTGCGCAACGTTGTTGCCATTGCTACAGGCATC
GTGGTGTACGCTCGTCTGTTGGTATGGCTTCATTGAGCTCCGGTTCCCAACGATCAAG
GCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCTCCGA
TCGTTGTGTCAGAAGTAAGTTGGCCGCAGTGTTATCACTCATGGTTATGGCAGCACTGCATA
ATTCTCTTACTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCA
AGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGG
GATAATACCGCGCCACATAGCAGAACTTTAAAGTGCTCATCATTGGAACGTTCTTCG
GGGCGAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTTCGATGTAACCCACTCG
TGCACCCAACTGATCTTCAGCATCTTTTACTTTACCAGCGTTTCTGGGTGAGCAAAAAC
AGGAAGGCAAAATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCA
TACTCTTCCTTTTTCAATATTATTGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATA
CATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAA
GTGCCACCTAAATTGTAAGCGTTAATATTTTGTAAATTCGCGTTAAATTTTTGTAAATC
AGCTCATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCCTTATAAATCAAAAGAATAGA
CCGAGATAGGGTTGAGTGTTGTTCCAGTTTGAACAAGAGTCCACTATTAAAGAACGTG
GACTCCAACGTCAAAGGGCGAAAAACCGTCTATCAGGGCGATGGCCCACTACGTGAAC
CATCACCTAATCAAGTTTTTTGGGGTTCGAGGTGCCGTAAAGCACTAAATCGGAACCCTA
AAGGGAGCCCCGATTTAGAGCTTGACGGGGAAAGCCAACCTGGCTTATCGAAATTAAT
ACGACTCACTATAGGGAGACCGGC

FIGURE 19

Schematic of pONY8.1Zhyb




Fig 20. Configuration of expression cassettes for RNAi





Fig 2. Aptazymes for regulated siRNA mediated gene silencing

A: design of expression cassette



B: structure of transcript






Fig 22. Hypoxically induced silencing of VEGF by siRNAs



In the above example the RNA polymerase III U6-RNA gene promoter is used to drive expression of an aptazyme-linked short hairpin against VEGF. Under hypoxic conditions expression from the hypoxic response element (HRE) is induced transcribing gene X which codes for a protein which is a ligand for the aptamer. Binding of the ligand to the aptazyme triggers catalysis and consequently gene silencing of vascular endothelial growth factor (VEGF)

Fig 23. Strategy for using RNA polymerase II promoters

In this case expression of the short hairpin is under the control of an RNA polymerase II promoter, CMV, Which also has two copies of Tet operator downstream providing an additional level of regulation.

Transcription is inhibited in the presence of the Tet repressor protein (in the absence of doxycycline) which may be expressed separately or from the same vector. The transcript is flanked by aptazymes which can be activated to cleave at sites designed to release the short hairpin such that it can initiate gene silencing of the target. It is necessary to use aptazymes rather than ribozymes as the latter would result in autologous cleavage of the lentiviral genome.



In the above configuration only expression of the antisense siRNA is under the control of aptazyme regulation. Again the target sequence is flanked by aptazymes which cleave at sites releasing the appropriate RNA sequence to form a duplex with the sense RNA which is constitutively expressed from the U6 promoter. Gene silencing of the target can therefore be switched on or off depending on the presence/absence of the aptazyme ligand.

Fig 24. Modulating expression of therapeutic sequences
A - Insulin







Fig 25. A Split intron strategy to avoid self-cleavage of RNA genome



The sequence coding for the aptazyme shown above would be split apart in the genome packaged by viral producer cells such that the region indicated in blue would not be present upstream of the sequence shown in black. Instead it will be present in the 3' LTR along with additional sequence comprising a splice donor (underlined). Upon reverse transcription this will be copied to the 5' LTR such that it is now upstream of a splice acceptor adjacent to the rest of the aptazyme sequence. Upon transcription the intron sequence, shown in red, will be spliced thereby forming a complete aptazyme. Therefore the aptazyme is only present in transduced cells preventing any prior cleavage of the vector genome which would lead to loss of it.


Fig 25B Vector configuration



Transduced

The vector depicted has an ElAV / MLV hybrid LTR. This also has a splice donor inserted downstream of the initiation of transcription and upstream of the ElAV repeat and which contains sequence of the 5' portion of the aptazyme. During reverse transcription the modified 3' LTR is copied to the 5' LTR. Following transcription and splicing the functional aptazyme is created. Activation of the aptazyme would cleave the transcript resulting in its degradation.

Fig 26 . 'Double hairpin' strategy to avoid self-cleavage of RNA genome



In the above example the 3' end of the U6 promoter has been modified to incorporate sequence which will base pair with the 5' region of helix I forming a hairpin which will prevent the aptazyme from adopting the configuration necessary for catalytic activity. This will only occur in the RNA genome and not in the transcript as initiation of transcription will be downstream of the sequence modified in the promoter.

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 July 2003 (10.07.2003)


PCT

(10) International Publication Number
WO 2003/056022 A3

- (51) International Patent Classification⁷: C12N 15/867, A01K 67/027
- (21) International Application Number: PCT/GB2002/005901
- (22) International Filing Date: 23 December 2002 (23.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0130797.4 21 December 2001 (21.12.2001) GB
0201140.1 18 January 2002 (18.01.2002) GB
10/082,122 26 February 2002 (26.02.2002) US
0211409.8 17 May 2002 (17.05.2002) GB
- (71) Applicant (for all designated States except US): OXFORD BIOMEDICA (UK) LIMITED [GB/GB]; Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).
- (72) Inventors; and
(75) Inventors/Applicants (for US only): RADCLIFFE, Philippa [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). MITROPHANOUS, Kyriacos [GR/GB]; c/o Oxford BioMedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). THEMIS, Michael [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).
- (74) Agents: MALLALIEU, Catherine, Louise et al.; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

[Continued on next page]

(54) Title: METHOD FOR PRODUCING A TRANSFENIC ORGANISM USING A LENTIVIRAL EXPRESSION VECTOR SUCH AS EIAV



(57) Abstract: A method of producing a transgenic cell comprising into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI). Also described is a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising a NOI capable of generating an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA, a micro-RNA or a group 1 intron. Also described is a viral vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises: (a) a second nucleotide sequence comprising an aptazyme; and (b) a third nucleotide sequence capable of generating a polynucleotide; wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK,

TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *with international search report*

(88) **Date of publication of the international search report:**

31 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 02/05901

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/867 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE, MEDLINE, PASCAL, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 6 312 683 B1 (CARROLL MILES WILLIAM ET AL) 6 November 2001 (2001-11-06) examples 4,5	1-6,15, 32,62
X	YAMADA K ET AL: "Functional correction of fanconi anemia group C hematopoietic cells by the use of a novel lentiviral vector." MOLECULAR THERAPY: THE JOURNAL OF THE AMERICAN SOCIETY OF GENE THERAPY. UNITED STATES APR 2001, vol. 3, no. 4, April 2001 (2001-04), pages 485-490, XP001147753 ISSN: 1525-0016 page 490, left-hand column, last paragraph; figure 2	1-6,15, 32,62

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

17 April 2003

Date of mailing of the international search report

10.09.03

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lanzrein, M

INTERNATIONAL SEARCH REPORT

International Application No
PC 1/ GB 02/05901

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TARANTAL A F ET AL: "Lentiviral vector gene transfer into fetal rhesus monkeys (Macaca mulatta): lung-targeting approaches."</p> <p>MOLECULAR THERAPY: THE JOURNAL OF THE AMERICAN SOCIETY OF GENE THERAPY. UNITED STATES DEC 2001, vol. 4, no. 6, December 2001 (2001-12), pages 614-621, XP002238838 ISSN: 1525-0016 figures 1,3,5; table 2</p> <p>-----</p> <p>Bathmore of relevance?</p>	<p>1-16, 32-39, 62-65</p>

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB 02/05901

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-16, 62 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-16 (completely); 32-39, 62-65 (partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 02 05901

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-16 (completely); 32-39, 62-65 (partially)

A method for producing a transgenic cell using a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI); the transgenic cell produced by said method; a transgenic organism which is generated from said transgenic cell.

2. Claims: 17-31 (completely); 32-39, 62-65 (partially)

A method for producing a transgenic cell by introducing into a non-dividing cell lentiviral expression vector comprising a nucleotide of interest (NOI); the transgenic cell produced by said method; a transgenic organism which is generated from said transgenic cell.

3. Claims: 40-61 (completely); 62-65 (partially)

A vector comprising an aptazyme operably linked to a sequence capable of generating a polynucleotide, wherein the aptazyme is activable to cleave a transcript such that said polynucleotide is generated; method for producing a transgenic cell using said vector; a transgenic organism generated from said transgenic cell.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 02/05901

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6312683	B1	06-11-2001	AU 760287 B2	08-05-2003
			AU 1773299 A	12-07-1999
			CA 2315269 A1	01-07-1999
			CN 1285000 T	21-02-2001
			EP 1042493 A1	11-10-2000
			GB 2356200 A ,B	16-05-2001
			WO 9932646 A1	01-07-1999
			GB 2334257 A ,B	18-08-1999
			GB 2345062 A ,B	28-06-2000
			JP 2002516065 T	04-06-2002

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 July 2003 (10.07.2003)

PCT


(10) International Publication Number
WO 2003/056022 A3

- (51) International Patent Classification⁷: **C12N 15/867**, (71) Applicant (for all designated States except US): **OXFORD BIOMEDICA (UK) LIMITED** [GB/GB]; Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).
- (21) International Application Number: **PCT/GB2002/005901** (72) Inventors; and (75) Inventors/Applicants (for US only): **RADCLIFFE, Philippa** [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). **MITROPHANOUS, Kyriacos** [GR/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB). **THEMIS, Michael** [GB/GB]; c/o Oxford Biomedica (UK) Limited, Medawar Centre, Robert Robinson Avenue, The Oxford Science Park, Oxford OX4 4GA (GB).
- (22) International Filing Date: **23 December 2002 (23.12.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
- | | | |
|------------|-------------------------------|----|
| 0130797.4 | 21 December 2001 (21.12.2001) | GB |
| 0201140.1 | 18 January 2002 (18.01.2002) | GB |
| 10/082,122 | 26 February 2002 (26.02.2002) | US |
| 0211409.8 | 17 May 2002 (17.05.2002) | GB |
- (74) Agents: **MALLALIEU, Catherine, Louise et al.**; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).

[Continued on next page]

(54) Title: METHOD FOR PRODUCING A TRANSGENIC ORGANISM USING A LENTIVIRAL EXPRESSION VECTOR SUCH AS EIAV

Schematic of pONY8.1Zhyb



(57) Abstract: A method of producing a transgenic cell comprising into a cell a non-primate lentiviral expression vector comprising a nucleotide of interest (NOI). Also described is a method of producing a transgenic cell comprising introducing into a cell a lentiviral expression vector comprising a NOI capable of generating an antisense oligonucleotide, a ribozyme, an siRNA, a short hairpin RNA, a micro-RNA, a micro-RNA or a group 1 intron. Also described is a viral vector comprising a first nucleotide sequence, wherein said first nucleotide sequence comprises: (a) a second nucleotide sequence comprising an aptazyme; and (b) a third nucleotide sequence capable of generating a polynucleotide; wherein (a) and (b) are operably linked and wherein the aptazyme is activatable to cleave a transcript of the first nucleotide sequence such that said polynucleotide is generated.



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report
— with amended claims

(88) **Date of publication of the international search report:**
31 December 2003

Date of publication of the amended claims: 12 February 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

AMENDED CLAIMS

[received by the International Bureau on 10 November 2003 (10.11.03);
new claims 66-70 added; remaining claims unchanged (1 page)]

66. A method of producing a transgenic cell comprising introducing into a cell a viral expression vector comprising a nucleotide of interest (NOI) which encodes and is capable of expressing a therapeutic protein.

5

67. A method of producing a transgenic cell comprising introducing into a cell a viral expression vector comprising an aptazyme capable of generating an siRNA, an antisense oligonucleotide, a ribozyme, a short hairpin RNA, a micro-RNA or a group I intron.

10

68. A transgenic cell produced by the method of claim 66 or 67.

69. A transgenic organism which is generated from or obtainable by generation from a transgenic cell according to claim 68, or from the method as defined in claim

15

66 or 67.

70. A transgenic egg derived from a transgenic organism according to claim 69.] *

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)